

Sepracor

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 June 2003 (19.06.2003)

PCT

(10) International Publication Number
WO 03/049736 A1

(51) International Patent Classification⁷: **A61K 31/445**,
31/454, 31/4545, C07D 211/54, 211/68, 401/02, 401/12

(21) International Application Number: PCT/US02/39651

(22) International Filing Date:
11 December 2002 (11.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/339,506 11 December 2001 (11.12.2001) US

(71) Applicant: **SEPRACOR, INC.** [US/US]; 84 Waterford
Drive, Marlborough, MA 01752-7010 (US).

(72) Inventors: **RADEKE, Heike**; 15 Seaver Farm Lane,
South Grafton, MA 01560 (US). **SHAO, Liming**; 158
South Great Road, Lincoln, MA 01773 (US).

(74) Agents: **GORDON, Dana, M.** et al.; Foley Hoag LLP,
Patent Group, 155 Seaport Boulevard, Boston, MA 02210
(US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/049736 A1

(54) Title: 4-SUBSTITUTED PIPERIDINES, AND METHODS OF USE THEREOF

(57) Abstract: One aspect of the present invention relates to heterocyclic compounds. A second aspect of the present invention relates to the use of the heterocyclic compounds as ligands for various mammalian cellular receptors, including dopamine, serotonin, or norepinephrine transporters. The compounds of the present invention will find use in the treatment of numerous ailments, conditions and diseases which afflict mammals, including but not limited to addiction, anxiety, depression, sexual dysfunction, hypertension, migraine, Alzheimer's disease, obesity, emesis, psychosis, schizophrenia, Parkinson's disease, inflammatory pain, neuropathic pain, Lesche-Nyhan disease, Wilson's disease, and Tourette's syndrome. An additional aspect of the present invention relates to the synthesis of combinatorial libraries of the heterocyclic compounds, and the screening of those libraries for biological activity, e.g., in assays based on dopamine transporters.

4-SUBSTITUTED PIPERIDINES, AND METHODS OF USE THEREOF

Background of the Invention

Dopamine, norepinephrine and serotonin are mammalian monoamine
5 neurotransmitters that play important roles in a wide variety of physiological processes. Therefore, compounds that selectively modulate the activity of these three neurotransmitters, either individually, in pairs, or as a group, promise to serve as agents effective in the treatment of a wide range of maladies, conditions and diseases that afflict mammals due to atypical activities of these neurotransmitters.

10 For example, depression is believed to result from dysfunction in the noradrenergic, dopaminergic, or serotonergic systems. Furthermore, the noradrenergic system appears to be associated with increased drive, whereas the serotonergic system relates more to changes in mood. Therefore, it is possible that the different symptoms of depression may benefit from drugs acting mainly on one or the other of these neurotransmitter systems. On the
15 other hand, a single compound that selectively affects both the noradrenergic and serotonergic systems should prove effective in the treatment of depression comprising symptoms related to dysfunction in both systems.

Dopamine plays a major role in addiction. Many of the concepts that apply to dopamine apply to other neurotransmitters as well. As a chemical messenger, dopamine is
20 similar to adrenaline. Dopamine affects brain processes that control movement, emotional response, and ability to experience pleasure and pain. Regulation of dopamine plays a crucial role in our mental and physical health. Neurons containing the neurotransmitter dopamine are clustered in the midbrain in an area called the substantia nigra. In Parkinson's disease, the dopamine-transmitting neurons in this area die. As a result, the brains of
25 people with Parkinson's disease contain almost no dopamine. To help relieve their symptoms, these patients are given L-DOPA, a drug that can be converted in the brain to dopamine.

Certain drugs are known as dopamine agonists. These drugs bind to dopamine receptors in place of dopamine and directly stimulate those receptors. Some dopamine
30 agonists are currently used to treat Parkinson's disease. These drugs can stimulate dopamine receptors even in someone without dopamine-secreting neurons. In contrast to dopamine agonists, dopamine antagonists are drugs that bind but don't stimulate dopamine receptors.

Antagonists can prevent or reverse the actions of dopamine by keeping dopamine from activating receptors.

Dopamine antagonists are traditionally used to treat schizophrenia and related mental disorders. A person with schizophrenia may have an overactive dopamine system.

5 Dopamine antagonists can help regulate this system by "turning down" dopamine activity.

Cocaine and other drugs of abuse can alter dopamine function. Such drugs may have very different actions. The specific action depends on which dopamine receptors and brain regions the drugs stimulate or block, and how well the compounds mimic dopamine. Drugs such as cocaine and amphetamine produce their effects by changing the flow of
10 neurotransmitters. These drugs are defined as indirect acting because they depend on the activity of neurons. In contrast, some drugs bypass neurotransmitters altogether and act directly on receptors.

Use of these two types of drugs can lead to very different results in treating the same disease. As mentioned earlier, people with Parkinson's disease lose neurons that contain
15 dopamine. To compensate for this loss, the body produces more dopamine receptors on other neurons. Indirect agonists are not very effective in treating the disease since they depend on the presence of dopamine neurons. In contrast, direct agonists are more effective because they stimulate dopamine receptors even when dopamine neurons are missing.

Certain drugs increase dopamine concentrations by preventing dopamine reuptake,
20 leaving more dopamine in the synapse. An example is the widely abused stimulant drug, cocaine. Another example is methylphenidate, used therapeutically to treat childhood hyperkinesia and symptoms of narcolepsy.

Sensitization or desensitization normally occurs with drug exposure. However, addiction or mental illness can tamper with the reuptake system. This disrupts the normal
25 levels of neurotransmitters in the brain and can lead to faulty desensitization or sensitization. If this happens in a region of the brain that serves emotion or motivation, the individual can suffer severe consequences. For example, cocaine prevents dopamine reuptake by binding to proteins that normally transport dopamine. Not only does cocaine "bully" dopamine out of the way, it also hangs on to the transport proteins much longer than
30 dopamine does. As a result, more dopamine remains to stimulate neurons, which causes a prolonged feelings of pleasure and excitement. Amphetamine also increases dopamine levels. Again, the result is over-stimulation of these pleasure-pathway nerves in the brain.

Dopamine activity is implicated in the reinforcing effects of cocaine, amphetamine and natural rewards. However, dopamine abnormalities are also believed to underlie some of the core attention deficits seen in acute schizophrenics.

Norepinephrine, also called noradrenaline, is a neurotransmitter that also acts as a hormone. As a neurotransmitter, norepinephrine helps to regulate arousal, dreaming, and moods. As a hormone, it acts to increase blood pressure, constrict blood vessels and increase heart rate - responses that occur when we feel stress.

Serotonin (5-hydroxytryptamine, 5-HT) is widely distributed in animals and plants, occurring in vertebrates, fruits, nuts, and venoms. A number of congeners of serotonin are also found in nature and have been shown to possess a variety of peripheral and central nervous system activities. Serotonin may be obtained from a variety of dietary sources; however, endogenous 5-HT is synthesized *in situ* from tryptophan through the actions of the enzymes tryptophan hydroxylase and aromatic L-amino acid decarboxylase. Both dietary and endogenous 5-HT are rapidly metabolized and inactivated by monoamine oxidase and aldehyde dehydrogenase to the major metabolite, 5-hydroxyindoleacetic acid (5-HIAA).

Serotonin is implicated in the etiology or treatment of various disorders, particularly those of the central nervous system, including anxiety, depression, obsessive-compulsive disorder, schizophrenia, stroke, obesity, pain, hypertension, vascular disorders, migraine, and nausea. Recently, understanding of the role of 5-HT in these and other disorders has advanced rapidly due to increasing understanding of the physiological role of various serotonin receptor subtypes.

It is currently estimated that up to 30% of clinically diagnosed cases of depression are resistant to all forms of drug therapy. To achieve an effective therapy for such patients, it is logical to develop drugs that possess reuptake inhibition profiles different from those of drugs currently available on the market. For example, the exact role of dopamine in depressive illness is far from clear; however, intervention in the dopamine system may hold promise for the treatment of a subset of major depression.

Summary of the Invention

One aspect of the present invention relates to heterocyclic compounds. A second aspect of the present invention relates to the use of the heterocyclic compounds as ligands for various mammalian cellular receptors, including dopamine, serotonin, or norepinephrine transporters. The compounds of the present invention will find use in the treatment of

numerous ailments, conditions and diseases which afflict mammals, including but not limited to addiction, anxiety, depression, sexual dysfunction, hypertension, migraine, Alzheimer's disease, obesity, emesis, psychosis, analgesia, schizophrenia, Parkinson's disease, restless leg syndrome, sleeping disorders, attention deficit hyperactivity disorder, irritable bowel syndrome, premature ejaculation, menstrual dysphoria syndrome, urinary incontinence, inflammatory pain, neuropathic pain, Lesche-Nyhan disease, Wilson's disease, and Tourette's syndrome. An additional aspect of the present invention relates to the synthesis of combinatorial libraries of the heterocyclic compounds, and the screening of those libraries for biological activity, e.g., in assays based on dopamine transporters.

10 ***Detailed Description of the Invention***

The present invention provides heterocyclic compounds, and combinatorial libraries thereof. Furthermore, the present invention provides heterocyclic compounds that are ligands for dopamine, serotonin, or norepinephrine receptors or transporters, and methods of use thereof for the treatment of numerous ailments, conditions and diseases which afflict mammals, including but not limited to addiction, anxiety, depression, sexual dysfunction, hypertension, migraine, Alzheimer's disease, obesity, emesis, psychosis, analgesia, schizophrenia, Parkinson's disease, restless leg syndrome, sleeping disorders, attention deficit hyperactivity disorder, irritable bowel syndrome, premature ejaculation, menstrual dysphoria syndrome, urinary incontinence, inflammatory pain, neuropathic pain, Lesche-Nyhan disease, Wilson's disease, and Tourette's syndrome. The present invention also relates to pharmaceutical formulations of the heterocyclic compounds.

In certain embodiments, compounds of the present invention are ligands for mammalian receptors for dopamine, norepinephrine, serotonin, any two of these three neurotransmitters or all of them. In certain embodiments, compounds of the present invention are ligands for mammalian transporters of dopamine, norepinephrine, serotonin, any two of these three neurotransmitters or all of them. In certain embodiments, compounds of the present invention are agonists of mammalian receptors for dopamine, norepinephrine, serotonin, any two of these three neurotransmitters or all of them. In certain embodiments, compounds of the present invention are antagonists or inverse agonists of mammalian receptors for dopamine, norepinephrine, serotonin, any two of these three neurotransmitters or all of them. In certain embodiments, compounds of the present invention are agonists of mammalian transporters of dopamine, norepinephrine, serotonin, any two of these three neurotransmitters or all of them. In certain embodiments,

compounds of the present invention are antagonists or inverse agonists of mammalian transporters of dopamine, norepinephrine, serotonin, any two of these three neurotransmitters or all of them.

In certain embodiments, compounds of the present invention are ligands for
5 mammalian dopamine receptors. In certain embodiments, compounds of the present invention are ligands for mammalian dopamine transporters. In certain embodiments, compounds of the present invention are agonists of mammalian dopamine receptors. In certain embodiments, compounds of the present invention are antagonists or inverse
10 agonists of mammalian dopamine receptors. In certain embodiments, compounds of the present invention are agonists of mammalian dopamine transporters. In certain embodiments, compounds of the present invention are antagonists or inverse agonists of mammalian dopamine transporters.

The mammalian dopamine receptor and transporter are members of a family of cell surface proteins that permit intracellular transduction of extracellular signals. Cell surface
15 proteins provide eukaryotic and prokaryotic cells a means to detect extracellular signals and transduce such signals intracellularly in a manner that ultimately results in a cellular response or a concerted tissue or organ response. Cell surface proteins, by intracellularly transmitting information regarding the extracellular environment via specific intracellular pathways induce an appropriate response to a particular stimulus. The response may be
20 immediate and transient, slow and sustained, or some mixture thereof. By virtue of an array of varied membrane surface proteins, eukaryotic cells are exquisitely sensitive to their environment.

Extracellular signal molecules, such as growth hormones, vasodilators and neurotransmitters, exert their effects, at least in part, via interaction with cell surface
25 proteins. For example, some extracellular signal molecules cause changes in transcription of target gene via changes in the levels of secondary messengers, such as cAMP. Other signals, indirectly alter gene expression by activating the expression of genes, such as immediate-early genes that encode regulatory proteins, which in turn activate expression of other genes that encode transcriptional regulatory proteins. For example, neuron gene
30 expression is modulated by numerous extracellular signals, including neurotransmitters and membrane electrical activity. Transsynaptic signals cause rapid responses in neurons that occur over a period of time ranging from milliseconds, such as the opening of ligand-gated channels, to seconds and minutes, such as second messenger-mediated events. Genes in

neural cells that are responsive to transsynaptic stimulation and membrane electrical activity, include genes, called immediate early genes, whose transcription is activated rapidly, within minutes, and transiently (see, e.g., Sheng et al. (1990) Neuron 4: 477-485), and genes whose expression requires protein synthesis and whose expression is induced or
5 altered over the course of hours.

Cell surface receptors and ion channels are among the cell surface proteins that respond to extracellular signals and initiate the events that lead to this varied gene expression and response. Ion channels and cell surface-localized receptors are ubiquitous and physiologically important cell surface membrane proteins. They play a central role in
10 regulating intracellular levels of various ions and chemicals, many of which are important for cell viability and function.

Cell surface-localized receptors are membrane spanning proteins that bind extracellular signalling molecules or changes in the extracellular environment and transmit the signal via signal transduction pathways to effect a cellular response. Cell surface
15 receptors bind circulating signal polypeptides, such as neurotransmitters, growth factors and hormones, as the initiating step in the induction of numerous intracellular pathways. Receptors are classified on the basis of the particular type of pathway that is induced. Included among these classes of receptors are those that bind growth factors and have intrinsic tyrosine kinase activity, such as the heparin binding growth factor (HBGF)
20 receptors, and those that couple to effector proteins through guanine nucleotide binding regulatory proteins, which are referred to as G protein coupled receptors and G proteins, respectively.

The G protein transmembrane signaling pathways consist of three proteins: receptors, G proteins and effectors. G proteins, which are the intermediaries in
25 transmembrane signaling pathways, are heterodimers and consist of alpha , beta and gamma subunits. Among the members of a family of G proteins the alpha subunits differ. Functions of G proteins are regulated by the cyclic association of GTP with the alpha subunit followed by hydrolysis of GTP to GDP and dissociation of GDP.

G protein coupled receptors are a diverse class of receptors that mediate signal
30 transduction by binding to G proteins. Signal transduction is initiated via ligand binding to the cell membrane receptor, which stimulates binding of the receptor to the G protein. The receptor G protein interaction releases GDP, which is specifically bound to the G protein, and permits the binding of GTP, which activates the G protein. Activated G protein

dissociates from the receptor and activates the effector protein, which regulates the intracellular levels of specific second messengers. Examples of such effector proteins include adenylyl cyclase, guanylyl cyclase, phospholipase C, and others.

G protein-coupled receptors, which are glycoproteins, are known to share certain structural similarities and homologies (see, e-g., Gilman, A.G., Ann. Rev. Biochem. 56: 615-649 (1987), Strader, C.D. et al. The FASEB Journal 3: 1825-1832 (1989), Kobilka, B.K., et al. Nature 329:75-79 (1985) and Young et al. Cell 45: 711-719 (1986)). Among the G protein-coupled receptors that have been identified and cloned are the substance P receptor, the angiotensin receptor, the alpha - and beta -adrenergic receptors and the serotonin receptors. G protein-coupled receptors share a conserved structural motif. The general and common structural features of the G protein-coupled receptors are the existence of seven hydrophobic stretches of about 20-25 amino acids each surrounded by eight hydrophilic regions of variable length. It has been postulated that each of the seven hydrophobic regions forms a transmembrane alpha helix and the intervening hydrophilic regions form alternately intracellularly and extracellularly exposed loops. The third cytosolic loop between transmembrane domains five and six is the intracellular domain responsible for the interaction with G proteins.

G protein-coupled receptors are known to be inducible. This inducibility was originally described in lower eukaryotes. For example, the cAMP receptor of the cellular slime mold, Dictyostelium, is induced during differentiation (Klein et al., Science 241: 1467-1472 (1988)). During the Dictyostelium discoideum differentiation pathway, cAMP, induces high level expression of its G protein-coupled receptor. This receptor transduces the signal to induce the expression of the other genes involved in chemotaxis, which permits multicellular aggregates to align, organize and form stalks (see, Firtel, R.A., et al. Cell 58: 235-239 (1989) and Devreotes, P., Science 245: 1054-1058 (1989)).

Definitions

For convenience, certain terms employed in the specification, examples, and appended claims are collected here.

The term "cell surface proteins" includes molecules that occur on the surface of cells, interact with the extracellular environment, and transmit or transduce information regarding the environment intracellularly.

The term "extracellular signals" includes a molecule or a change in the environment that is transduced intracellularly via cell surface proteins that interact, directly or indirectly,

with the signal. An extracellular signal is any compound or substance that in some manner specifically alters the activity of a cell surface protein. Examples of such signals include, but are not limited to, molecules such as acetylcholine, growth factors, hormones and other mitogenic substances, such as phorbol myristate acetate (PMA), that bind to cell surface
5 receptors and ion channels and modulate the activity of such receptors and channels. Extracellular signals also includes as yet unidentified substances that modulate the activity of a cell surface protein and thereby affect intracellular functions and that are potential pharmacological agents that may be used to treat specific diseases by modulating the activity of specific cell surface receptors.

10 The term "ED₅₀" means the dose of a drug which produces 50% of its maximum response or effect. Alternatively, the dose which produces a pre-determined response in 50% of test subjects or preparations.

The term "LD₅₀" means the dose of a drug which is lethal in 50% of test subjects.

The term "therapeutic index" refers to the therapeutic index of a drug defined as
15 LD₅₀/ED₅₀.

The term "structure-activity relationship (SAR)" refers to the way in which altering the molecular structure of drugs alters their interaction with a receptor, enzyme, etc.

The term "agonist" refers to a compound that mimics the action of natural transmitter or, when the natural transmitter is not known, causes changes at the receptor
20 complex in the absence of other receptor ligands.

The term "antagonist" refers to a compound that binds to a receptor site, but does not cause any physiological changes unless another receptor ligand is present.

The term "inverse agonist" refers to a compound that binds to a constitutively active receptor site and reduces its physiological function.

25 The term "competitive antagonist" refers to an antagonist, the effects of which can be overcome by increased concentration of an agonist.

The term "partial agonist" refers to a compound that binds to a receptor site but does not produce the maximal effect regardless of its concentration.

The term "ligand" refers to a compound that binds at the receptor site.

30 The term "heteroatom" as used herein means an atom of any element other than carbon or hydrogen. Preferred heteroatoms are boron, nitrogen, oxygen, phosphorus, sulfur and selenium.

The term "electron-withdrawing group" is recognized in the art, and denotes the tendency of a substituent to attract valence electrons from neighboring atoms, i.e., the substituent is electronegative with respect to neighboring atoms. A quantification of the level of electron-withdrawing capability is given by the Hammett sigma (σ) constant. This well known constant is described in many references, for instance, J. March, Advanced Organic Chemistry, McGraw Hill Book Company, New York, (1977 edition) pp. 251-259. The Hammett constant values are generally negative for electron donating groups ($\sigma[P] = -0.66$ for NH_2) and positive for electron withdrawing groups ($\sigma[P] = 0.78$ for a nitro group), $\sigma[P]$ indicating para substitution. Exemplary electron-withdrawing groups include nitro, acyl, formyl, sulfonyl, trifluoromethyl, cyano, chloride, and the like. Exemplary electron-donating groups include amino, methoxy, and the like.

The term "alkyl" refers to the radical of saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In preferred embodiments, a straight chain or branched chain alkyl has 30 or fewer carbon atoms in its backbone (e.g., $\text{C}_1\text{-C}_{30}$ for straight chain, $\text{C}_3\text{-C}_{30}$ for branched chain), and more preferably 20 or fewer. Likewise, preferred cycloalkyls have from 3-10 carbon atoms in their ring structure, and more preferably have 5, 6 or 7 carbons in the ring structure.

Unless the number of carbons is otherwise specified, "lower alkyl" as used herein means an alkyl group, as defined above, but having from one to ten carbons, more preferably from one to six carbon atoms in its backbone structure. Likewise, "lower alkenyl" and "lower alkynyl" have similar chain lengths. Preferred alkyl groups are lower alkyls. In preferred embodiments, a substituent designated herein as alkyl is a lower alkyl.

The term "aralkyl", as used herein, refers to an alkyl group substituted with an aryl group (e.g., an aromatic or heteroaromatic group).

The terms "alkenyl" and "alkynyl" refer to unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond respectively.

The term "aryl" as used herein includes 5-, 6- and 7-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, pyrrole, furan, thiophene, imidazole, oxazole, thiazole, triazole, pyrazole, pyridine, pyrazine, pyridazine and pyrimidine, and the like. Those aryl groups having heteroatoms in the ring

structure may also be referred to as "aryl heterocycles" or "heteroaromatics." The aromatic ring can be substituted at one or more ring positions with such substituents as described above, for example, halogen, azide, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, alkoxy, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, sulfonamido, ketone, aldehyde, ester, heterocyclyl, aromatic or heteroaromatic moieties, -CF₃, -CN, or the like. The term "aryl" also includes polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings (the rings are "fused rings") wherein at least one of the rings is aromatic, e.g., the other cyclic rings can be cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls.

The terms *ortho*, *meta* and *para* apply to 1,2-, 1,3- and 1,4-disubstituted benzenes, respectively. For example, the names 1,2-dimethylbenzene and *ortho*-dimethylbenzene are synonymous.


The terms "heterocyclyl" or "heterocyclic group" refer to 3- to 10-membered ring structures, more preferably 3- to 7-membered rings, whose ring structures include one to four heteroatoms. Heterocycles can also be polycycles. Heterocyclyl groups include, for example, thiophene, thianthrene, furan, pyran, isobenzofuran, chromene, xanthene, phenoxathiin, pyrrole, imidazole, pyrazole, isothiazole, isoxazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, pyrimidine, phenanthroline, phenazine, phenarsazine, phenothiazine, furazan, phenoxazine, pyrrolidine, oxolane, thiolane, oxazole, piperidine, piperazine, morpholine, lactones, lactams such as azetidinones and pyrrolidinones, sultams, sultones, and the like. The heterocyclic ring can be substituted at one or more positions with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like.

The terms "polycyclyl" or "polycyclic group" refer to two or more rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more carbons are common to two adjoining rings, e.g., the rings are "fused rings". Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the

polycycle can be substituted with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -
 5 CF₃, -CN, or the like.


As used herein, the term "nitro" means -NO₂; the term "halogen" designates -F, -Cl, -Br or -I; the term "sulfhydryl" means -SH; the term "hydroxyl" means -OH; and the term "sulfonyl" means -SO₂-.

The terms "amine" and "amino" are art-recognized and refer to both unsubstituted
 10 and substituted amines, e.g., a moiety that can be represented by the general formula:




wherein R₉, R₁₀ and R'₁₀ each independently represent a group permitted by the rules of valence.

The term "acylamino" is art-recognized and refers to a moiety that can be
 15 represented by the general formula:



wherein R₉ represents a group permitted by the rules of valence, and R'₁₁ represents hydrogen, alkyl, cycloalkyl, alkenyl, aryl, heteroaryl, aralkyl, or heteroaralkyl.

The term "amido" is art recognized as an amino-substituted carbonyl and includes a
 20 moiety that can be represented by the general formula:




wherein R₉, R₁₀ are as defined above. Preferred embodiments of the amide will not include imides which may be unstable.

The term "alkylthio" refers to an alkyl group, as defined above, having a sulfur
 25 radical attached thereto. In preferred embodiments, the "alkylthio" moiety is represented by

one of -S-alkyl, -S-alkenyl, -S-alkynyl, and -S-(CH₂)_m-R₈, wherein m is an integer less than or equal to ten, and R₈ is alkyl, cycloalkyl, alkenyl, aryl, or heteroaryl. Representative alkylthio groups include methylthio, ethyl thio, and the like.

The term "carbonyl" is art recognized and includes such moieties as can be
 5 represented by the general formula:




wherein X is a bond or represents an oxygen or a sulfur, and R₁₁ represents a hydrogen, an alkyl, an alkenyl, -(CH₂)_m-R₈ or a pharmaceutically acceptable salt, R'₁₁ represents a hydrogen, an alkyl, an alkenyl or -(CH₂)_m-R₈, where m and R₈ are as defined above.

10 Where X is an oxygen and R₁₁ or R'₁₁ is not hydrogen, the formula represents an "ester". Where X is an oxygen, and R₁₁ is as defined above, the moiety is referred to herein as a carboxyl group, and particularly when R₁₁ is a hydrogen, the formula represents a "carboxylic acid". Where X is an oxygen, and R'₁₁ is hydrogen, the formula represents a "formate". In general, where the oxygen atom of the above formula is replaced by sulfur,
 15 the formula represents a "thiolcarbonyl" group. Where X is a sulfur and R₁₁ or R'₁₁ is not hydrogen, the formula represents a "thiolester." Where X is a sulfur and R₁₁ is hydrogen, the formula represents a "thiolcarboxylic acid." Where X is a sulfur and R'₁₁ is hydrogen, the formula represents a "thiolformate." On the other hand, where X is a bond, and R₁₁ is not hydrogen, the above formula represents a "ketone" group. Where X is a bond, and R₁₁
 20 is hydrogen, the above formula represents an "aldehyde" group.

The terms "alkoxyl" or "alkoxy" as used herein refers to an alkyl group, as defined above, having an oxygen radical attached thereto. Representative alkoxyl groups include methoxy, ethoxy, propyloxy, tert-butoxy and the like. An "ether" is two hydrocarbons covalently linked by an oxygen. Accordingly, the substituent of an alkyl that renders that
 25 alkyl an ether is or resembles an alkoxyl, such as can be represented by one of -O-alkyl, -O-alkenyl, -O-alkynyl, -O-(CH₂)_m-R₈, where m and R₈ are described above.

The term "sulfonate" is art recognized and includes a moiety that can be represented by the general formula:




in which R₄₁ is an electron pair, hydrogen, alkyl, cycloalkyl, or aryl.

The terms triflyl, tosyl, mesyl, and nonafllyl are art-recognized and refer to trifluoromethanesulfonyl, *p*-toluenesulfonyl, methanesulfonyl, and

5 nonafluorobutanesulfonyl groups, respectively. The terms triflate, tosylate, mesylate, and nonaflate are art-recognized and refer to trifluoromethanesulfonate ester, *p*-toluenesulfonate ester, methanesulfonate ester, and nonafluorobutanesulfonate ester functional groups and molecules that contain said groups, respectively.


The abbreviations Me, Et, Ph, Tf, Nf, Ts, Ms represent methyl, ethyl, phenyl, 10 trifluoromethanesulfonyl, nonafluorobutanesulfonyl, *p*-toluenesulfonyl and methanesulfonyl, respectively. A more comprehensive list of the abbreviations utilized by organic chemists of ordinary skill in the art appears in the first issue of each volume of the *Journal of Organic Chemistry*; this list is typically presented in a table entitled Standard List of Abbreviations. The abbreviations contained in said list, and all abbreviations 15 utilized by organic chemists of ordinary skill in the art are hereby incorporated by reference.

The term "sulfate" is art recognized and includes a moiety that can be represented by the general formula:




20 in which R₄₁ is as defined above.


The term "sulfonylamino" is art recognized and includes a moiety that can be represented by the general formula:



The term "sulfamoyl" is art-recognized and includes a moiety that can be 25 represented by the general formula:




The term "sulfonyl", as used herein, refers to a moiety that can be represented by the general formula:



- 5 in which R₄₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl.

The term "sulfoxido" as used herein, refers to a moiety that can be represented by the general formula:



- 10 in which R₄₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aralkyl, or aryl.

A "selenoalkyl" refers to an alkyl group having a substituted seleno group attached thereto. Exemplary "selenoethers" which may be substituted on the alkyl are selected from one of -Se-alkyl, -Se-alkenyl, -Se-alkynyl, and -Se-(CH₂)_m-R₈, m and R₈ being defined

- 15 above.

Analogous substitutions can be made to alkenyl and alkynyl groups to produce, for example, aminoalkenyls, aminoalkynyls, amidoalkenyls, amidoalkynyls, iminoalkenyls, iminoalkynyls, thioalkenyls, thioalkynyls, carbonyl-substituted alkenyls or alkynyls.

- 20 As used herein, the definition of each expression, e.g. alkyl, m, n, etc., when it occurs more than once in any structure, is intended to be independent of its definition elsewhere in the same structure.

- It will be understood that "substitution" or "substituted with" includes the implicit proviso that such substitution is in accordance with permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, e.g., which
25 does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc.

As used herein, the term "substituted" is contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic substituents of organic compounds. Illustrative substituents include, for example, those described herein above. The permissible substituents can be one or more and the same or different for appropriate organic compounds. For purposes of this invention, the heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valences of the heteroatoms. This invention is not intended to be limited in any manner by the permissible substituents of organic compounds.

The phrase "protecting group" as used herein means temporary substituents which protect a potentially reactive functional group from undesired chemical transformations. Examples of such protecting groups include esters of carboxylic acids, silyl ethers of alcohols, and acetals and ketals of aldehydes and ketones, respectively. The field of protecting group chemistry has been reviewed (Greene, T.W.; Wuts, P.G.M. *Protective Groups in Organic Synthesis*, 2nd ed.; Wiley: New York, 1991).

Certain compounds of the present invention may exist in particular geometric or stereoisomeric forms. The present invention contemplates all such compounds, including *cis*- and *trans*-isomers, *R*- and *S*-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.


If, for instance, a particular enantiomer of a compound of the present invention is desired, it may be prepared by asymmetric synthesis, or by derivation with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide the pure desired enantiomers. Alternatively, where the molecule contains a basic functional group, such as amino, or an acidic functional group, such as carboxyl, diastereomeric salts are formed with an appropriate optically-active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers. Further, mixtures of stereoisomers may be resolved using chiral chromatographic means.

Contemplated equivalents of the compounds described above include compounds which otherwise correspond thereto, and which have the same general properties thereof, wherein one or more simple variations of substituents are made which do not adversely affect the efficacy of the compound in binding to monoamine transporters. In general, the compounds of the present invention may be prepared by the methods illustrated in the general reaction schemes as, for example, described below, or by modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants which are in themselves known, but are not mentioned here.

For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 67th Ed., 1986-87, inside cover.

Compounds of the Invention

In certain embodiments, a compound of the present invention is represented by A:



A

wherein

X represents O, S, SO, SO₂, NR₇, NC(O)R₇, NC(O)OR₇, or NS(O)₂R₇;

m is 0, 1, 2, 3, or 4;

y is 0, 1, or 2;

z is 0, 1, or 2;

R₄ represents H, alkyl, cycloalkyl, alkenyl, aryl, or heteroaryl;

R_5 and R_6 are selected independently for each occurrence from the group consisting of H, alkyl, aryl, heteroaryl, F, OR_{11} , and $OC(O)R_{11}$; or an instance of CR_5R_6 taken together is C(O);

R_7 represents alkyl, cycloalkyl, aryl, heteroaryl, aralkyl, or heteroaralkyl;

5 R_8 and R_9 are selected independently for each occurrence from the group consisting of H, alkyl, aryl, heteroaryl, F, OR_{11} , and $OC(O)R_{11}$; or an instance of CR_8R_9 taken together is C(O);

R_{10} represents H, alkyl, cycloalkyl, alkenyl, aryl, or heteroaryl;

R_{11} represents H, alkyl, cycloalkyl, alkenyl, aryl, or heteroaryl;

10 a covalent bond may connect R_4 and an instance of R_5 or R_6 ;

a covalent bond may connect R_{10} and an instance of R_8 or R_9 ;

any two instances of R_5 and R_6 may be connected through a covalent bond;

any two geminal or vicinal instances of R_8 and R_9 may be connected through a covalent bond; and

15 the stereochemical configuration at any stereocenter of a compound represented by A is R, S, or a mixture of these configurations.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O.

20 In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein m is 2.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein y is 0.

25 In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein y is 1.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein z is 0.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein z is 1.

30 In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein R_4 is aryl.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein R_4 is 4-chlorophenyl.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl.

In certain embodiments, the compounds of the present invention are represented by
5 A and the attendant definitions, wherein R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein R₁₀ is aryl.

In certain embodiments, the compounds of the present invention are represented by
10 A and the attendant definitions, wherein R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; and R₄ is aryl.

In certain embodiments, the compounds of the present invention are represented by
15 A and the attendant definitions, wherein X is O; R₄ is aryl; and R₁₀ is aryl.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; R₄ is aryl; R₁₀ is aryl; and R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl.

In certain embodiments, the compounds of the present invention are represented by
20 A and the attendant definitions, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; and R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl.

In certain embodiments, the compounds of the present invention are represented by
25 A and the attendant definitions, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; and m is 2.

In certain embodiments, the compounds of the present invention are represented by
30 A and the attendant definitions, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; and y is 0.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; and y is 1.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; and z is 0.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; and z is 1.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; R₄ is 4-chlorophenyl; and R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; R₄ is 4-chlorophenyl; R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; and R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; R₄ is 4-chlorophenyl; R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; and m is 2.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; R₄ is 4-chlorophenyl; R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are

selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; y is 1; and z is 1.

In certain embodiments, the compounds of the present invention are represented by A and the attendant definitions, wherein X is O; R₄ is 4-chlorophenyl; R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl; R₅ and R₆ are selected
5 independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; y is 1; and z is 0.

10 In assays based on mammalian dopamine, serotonin, or norepinephrine receptors or transporters, certain compounds according to structure A have EC₅₀ values less than 1 μ M, more preferably less than 100 nM, and most preferably less than 10 nM.

In assays based on mammalian dopamine receptors or transporters, certain compounds according to structure A have EC₅₀ values less than 1 μ M, more preferably less
15 than 100 nM, and most preferably less than 10 nM.

In assays based on mammalian dopamine, serotonin, or norepinephrine receptors or transporters, certain compounds according to structure A have IC₅₀ values less than 1 μ M, more preferably less than 100 nM, and most preferably less than 10 nM.

In assays based on mammalian dopamine receptors or transporters, certain
20 compounds according to structure A have IC₅₀ values less than 1 μ M, more preferably less than 100 nM, and most preferably less than 10 nM.

In certain embodiments, compounds according to structure A are effective in the treatment of mammals suffering from addiction, anxiety, depression, sexual dysfunction, hypertension, migraine, Alzheimer's disease, obesity, emesis, psychosis, analgesia,
25 schizophrenia, Parkinson's disease, restless leg syndrome, sleeping disorders, attention deficit hyperactivity disorder, irritable bowel syndrome, premature ejaculation, menstrual dysphoria syndrome, urinary incontinence, inflammatory pain, neuropathic pain, Lesche-Nyhane disease, Wilson's disease, or Tourette's syndrome.

30 In certain embodiments, the present invention relates to a compound represented by any of the structures outlined above, wherein said compound is a single stereoisomer.

In certain embodiments, the present invention relates to a formulation, comprising a compound represented by any of the structures outlined above; and a pharmaceutically acceptable excipient.

In certain embodiments, the present invention relates to ligands for receptors or
5 transporters of dopamine, serotonin, or norepinephrine, wherein the ligands are represented by any of the structures outlined above, and any of the sets of definitions associated with one of those structures. In certain embodiments, the ligands of the present invention are antagonists or agonists of receptors or transporters of dopamine, serotonin, or
10 norepinephrine. In any event, the ligands of the present invention preferably exert their effect on the dopamine, serotonin, or norepinephrine receptors or transporters at a concentration less than about 1 micromolar, more preferably at a concentration less than about 100 nanomolar, and most preferably at a concentration less than 10 nanomolar.

In certain embodiments, the selectivity of a ligand for dopamine receptors or transporters renders that ligand an effective therapeutic agent for an acute or chronic
15 ailment, disease or malady. In certain embodiments, the selectivity of a ligand for dopamine receptors or transporters consists of a binding affinity for dopamine receptors or transporters at least a factor of ten greater than its binding affinity for receptors or transporters of other neurotransmitters. In certain embodiments, the selectivity of a ligand for dopamine receptors or transporters consists of a binding affinity for dopamine receptors
20 or transporters at least a factor of one hundred greater than its binding affinity for receptors or transporters of other neurotransmitters. In certain embodiments, the selectivity of a ligand for dopamine receptors or transporters consists of a binding affinity for dopamine receptors or transporters at least a factor of one thousand greater than its binding affinity for receptors or transporters of other neurotransmitters.

25 The present invention contemplates pharmaceutical formulations of the ligands of the present invention. In certain embodiments, the pharmaceutical formulations will comprise ligands of the present invention that selectively effect dopamine receptors or transporters, and thereby have a therapeutic effect on an acute or chronic ailment, disease or malady that is at least in part due to biochemical or physiological processes associated with
30 dopamine receptors or transporters. The *Background of the Invention* (see above) teaches examples of acute or chronic ailments, diseases or maladies that are caused or exacerbated by biochemical or physiological processes associated with dopamine receptors or transporters. One of ordinary skill in the art will be able to accumulate, by reference to the

scientific literature, a more comprehensive list of acute or chronic ailments, diseases or maladies that are caused or exacerbated by biochemical or physiological processes associated with dopamine receptors or transporters. The present invention contemplates pharmaceutical formulations of ligands of the present invention that will be of medicinal value against the aforementioned acute or chronic ailments, diseases or maladies.

Biochemical Activity at Cellular Receptors, and Assays to Detect That Activity

Assaying processes are well known in the art in which a reagent is added to a sample, and measurements of the sample and reagent are made to identify sample attributes stimulated by the reagent. For example, one such assay process concerns determining in a chromogenic assay the amount of an enzyme present in a biological sample or solution. Such assays are based on the development of a colored product in the reaction solution. The reaction develops as the enzyme catalyzes the conversion of a colorless chromogenic substrate to a colored product.

Another assay useful in the present invention concerns determining the ability of a ligand to bind to a biological receptor utilizing a technique well known in the art referred to as a radioligand binding assay. This assay accurately determines the specific binding of a radioligand to a targeted receptor through the delineation of its total and nonspecific binding components. Total binding is defined as the amount of radioligand that remains following the rapid separation of the radioligand bound in a receptor preparation (cell homogenates or recombinant receptors) from that which is unbound. The nonspecific binding component is defined as the amount of radioligand that remains following separation of the reaction mixture consisting of receptor, radioligand and an excess of unlabeled ligand. Under this condition, the only radioligand that remains represents that which is bound to components other than receptor. The specific radioligand bound is determined by subtracting the nonspecific from total radioactivity bound. For a specific example of radioligand binding assay for μ -opioid receptor, see Wang, J. B. et al. *FEBS Letters* 1994, 338, 217.

Assays useful in the present invention concern determining the activity of receptors the activation of which initiates subsequent intracellular events in which intracellular stores of calcium ions are released for use as a second messenger. Activation of some G-protein-coupled receptors stimulates the formation of inositol triphosphate (IP₃, a G-protein-coupled receptor second messenger) through phospholipase C-mediated hydrolysis of

phosphatidylinositol, Berridge and Irvine (1984). Nature 312:315-21. IP3 in turn stimulates the release of intracellular calcium ion stores.

A change in cytoplasmic calcium ion levels caused by release of calcium ions from intracellular stores is used to determine G-protein-coupled receptor function. This is another
5 type of indirect assay. Among G-protein-coupled receptors are muscarinic acetylcholine receptors (mAChR), adrenergic receptors, sigma receptors, serotonin receptors, dopamine receptors, angiotensin receptors, adenosine receptors, bradykinin receptors, metabotropic excitatory amino acid receptors and the like. Cells expressing such G-protein-coupled
10 receptors may exhibit increased cytoplasmic calcium levels as a result of contribution from both intracellular stores and via activation of ion channels, in which case it may be desirable although not necessary to conduct such assays in calcium-free buffer, optionally supplemented with a chelating agent such as EGTA, to distinguish fluorescence response resulting from calcium release from internal stores. Another type of indirect assay involves
15 determining the activity of receptors which, when activated, result in a change in the level of intracellular cyclic nucleotides, e.g., cAMP, cGMP. For example, activation of some dopamine, serotonin, metabotropic glutamate receptors and muscarinic acetylcholine receptors results in a decrease in the cAMP or cGMP levels of the cytoplasm.

Furthermore, there are cyclic nucleotide-gated ion channels, e.g., rod photoreceptor cell channels and olfactory neuron channels [see, Altenhofen, W. et al. (1991) Proc. Natl.
20 Acad. Sci U.S.A. 88:9868-9872 and Dhallan et al. (1990) Nature 347:184-187] that are permeable to cations upon activation by binding of cAMP or cGMP. A change in cytoplasmic ion levels caused by a change in the amount of cyclic nucleotide activation of photo-receptor or olfactory neuron channels is used to determine function of receptors that cause a change in cAMP or cGMP levels when activated. In cases where activation of the
25 receptor results in a decrease in cyclic nucleotide levels, it may be preferable to expose the cells to agents that increase intracellular cyclic nucleotide levels, e.g., forskolin, prior to adding a receptor-activating compound to the cells in the assay. Cell for this type of assay can be made by co-transfection of a host cell with DNA encoding a cyclic nucleotide-gated ion channel and a DNA encoding a receptor (e.g., certain metabotropic glutamate receptors,
30 muscarinic acetylcholine receptors, dopamine receptors, serotonin receptors and the like, which, when activated, causes a change in cyclic nucleotide levels in the cytoplasm.

Any cell expressing a receptor protein which is capable, upon activation, of directly increasing the intracellular concentration of calcium, such as by opening gated calcium

channels, or indirectly affecting the concentration of intracellular calcium as by causing initiation of a reaction which utilizes Ca^{2+} as a second messenger (e.g., G-protein-coupled receptors), may form the basis of an assay. Cells endogenously expressing such receptors or ion channels and cells which may be transfected with a suitable vector

5 encoding one or more such cell surface proteins are known to those of skill in the art or may be identified by those of skill in the art. Although essentially any cell which expresses endogenous ion channel and/or receptor activity may be used, it is preferred to use cells transformed or transfected with heterologous DNAs encoding such ion channels and/or receptors so as to express predominantly a single type of ion channel or receptor. Many

10 cells that may be genetically engineered to express a heterologous cell surface protein are known. Such cells include, but are not limited to, baby hamster kidney (BHK) cells (ATCC No. CCL10), mouse L cells (ATCC No. CCL1.3), DG44 cells [see, Chasin (1986) Cell. Molec. Genet. 12:555] human embryonic kidney (HEK) cells (ATCC No. CRL1573), Chinese hamster ovary (CHO) cells (ATCC Nos. CRL9618, CCL61, CRL9096), PC12 cells

15 (ATCC No. CRL1721) and COS-7 cells (ATCC No. CRL1651). Preferred cells for heterologous cell surface protein expression are those that can be readily and efficiently transfected. Preferred cells include HEK 293 cells, such as those described in U.S. Pat. No. 5,024,939.

Any compound which is known to activate ion channels or receptors of interest may

20 be used to initiate an assay. Choosing an appropriate ion channel- or receptor-activating reagent depending on the ion channel or receptor of interest is within the skill of the art. Direct depolarization of the cell membrane to determine calcium channel activity may be accomplished by adding a potassium salt solution having a concentration of potassium ions such that the final concentration of potassium ions in the cell-containing well is in the range

25 of about 50-150 mM (e.g., 50 mM KCl). With respect to ligand-gated receptors and ligand-gated ion channels, ligands are known which have affinity for and activate such receptors. For example, nicotinic acetylcholine receptors are known to be activated by nicotine or acetylcholine; similarly, muscarinic and acetylcholine receptors may be activated by addition of muscarine or carbamylcholine.

30 Agonist assays may be carried out on cells known to possess ion channels and/or receptors to determine what effect, if any, a compound has on activation or potentiation of ion channels or receptors of interest. Agonist assays also may be carried out using a reagent

known to possess ion channel- or receptor-activating capacity to determine whether a cell expresses the respective functional ion channel or receptor of interest.

Contacting a functional receptor or ion channel with agonist typically activates a transient reaction; and prolonged exposure to an agonist may desensitize the receptor or ion channel to subsequent activation. Thus, in general, assays for determining ion channel or receptor function should be initiated by addition of agonist (i.e., in a reagent solution used to initiate the reaction). The potency of a compound having agonist activity is determined by the detected change in some observable in the cells (typically an increase, although activation of certain receptors causes a decrease) as compared to the level of the observable in either the same cell, or substantially identical cell, which is treated substantially identically except that reagent lacking the agonist (i.e., control) is added to the well. Where an agonist assay is performed to test whether or not a cell expresses the functional receptor or ion channel of interest, known agonist is added to test-cell-containing wells and to wells containing control cells (substantially identical cell that lacks the specific receptors or ion channels) and the levels of observable are compared. Depending on the assay, cells lacking the ion channel and/or receptor of interest should exhibit substantially no increase in observable in response to the known agonist. A substantially identical cell may be derived from the same cells from which recombinant cells are prepared but which have not been modified by introduction of heterologous DNA. Alternatively, it may be a cell in which the specific receptors or ion channels are removed. Any statistically or otherwise significant difference in the level of observable indicates that the test compound has in some manner altered the activity of the specific receptor or ion channel or that the test cell possesses the specific functional receptor or ion channel.

In an example of drug screening assays for identifying compounds which have the ability to modulate ion channels or receptors of interest, individual wells (or duplicate wells, etc.) contain a distinct cell type, or distinct recombinant cell line expressing a homogeneous population of a receptor or ion channel of interest, so that the compound having unidentified activity may be screened to determine whether it possesses modulatory activity with respect to one or more of a variety of functional ion channels or receptors. It is also contemplated that each of the individual wells may contain the same cell type so that multiple compounds (obtained from different reagent sources in the apparatus or contained within different wells) can be screened and compared for modulating activity with respect to one particular receptor or ion channel type.

Antagonist assays, including drug screening assays, may be carried out by incubating cells having functional ion channels and/or receptors in the presence and absence of one or more compounds, added to the solution bathing the cells in the respective wells of the microtiter plate for an amount of time sufficient (to the extent that the
5 compound has affinity for the ion channel and/or receptor of interest) for the compound(s) to bind to the receptors and/or ion channels, then activating the ion channels or receptors by addition of known agonist, and measuring the level of observable in the cells as compared to the level of observable in either the same cell, or substantially identical cell, in the absence of the putative antagonist.

10 The assays are thus useful for rapidly screening compounds to identify those that modulate any receptor or ion channel in a cell. In particular, assays can be used to test functional ligand-receptor or ligand-ion channel interactions for cell receptors including ligand-gated ion channels, voltage-gated ion channels, G-protein-coupled receptors and growth factor receptors.

15 Those of ordinary skill in the art will recognize that assays may encompass measuring a detectable change of a solution as a consequence of a cellular event which allows a compound, capable of differential characteristics, to change its characteristics in response to the cellular event. By selecting a particular compound which is capable of differential characteristics upon the occurrence of a cellular event, various assays may be
20 performed. For example, assays for determining the capacity of a compound to induce cell injury or cell death may be carried out by loading the cells with a pH-sensitive fluorescent indicator such as BCECF (Molecular Probes, Inc., Eugene, Oreg. 97402, Catalog #B1150) and measuring cell injury or cell death as a function of changing fluorescence over time.

In a further example of useful assays, the function of receptors whose activation
25 results in a change in the cyclic nucleotide levels of the cytoplasm may be directly determined in assays of cells that express such receptors and that have been injected with a fluorescent compound that changes fluorescence upon binding cAMP. The fluorescent compound comprises cAMP-dependent-protein kinase in which the catalytic and regulatory subunits are each labelled with a different fluorescent-dye [Adams et al. (1991) Nature
30 349:694-697]. When cAMP binds to the regulatory subunits, the fluorescence emission spectrum changes; this change can be used as an indication of a change in cAMP concentration.

The function of certain neurotransmitter transporters which are present at the synaptic cleft at the junction between two neurons may be determined by the development of fluorescence in the cytoplasm of such neurons when conjugates of an amine acid and fluorescent indicator (wherein the fluorescent indicator of the conjugate is an
5 acetoxymethyl ester derivative e.g., 5-(aminoacetamido)fluorescein; Molecular Probes, Catalog #A1363) are transported by the neurotransmitter transporter into the cytoplasm of the cell where the ester group is cleaved by esterase activity and the conjugate becomes fluorescent.

In practicing an assay of this type, a reporter gene construct is inserted into an
10 eukaryotic cell to produce a recombinant cell which has present on its surface a cell surface protein of a specific type. The cell surface receptor may be endogenously expressed or it may be expressed from a heterologous gene that has been introduced into the cell. Methods for introducing heterologous DNA into eukaryotic cells are well known in the art and any such method may be used. In addition, DNA encoding various cell surface proteins is
15 known to those of skill in the art or it may be cloned by any method known to those of skill in the art.

The recombinant cell is contacted with a test compound and the level of reporter gene expression is measured. The contacting may be effected in any vehicle and the testing may be by any means using any protocols, such as serial dilution, for assessing specific
20 molecular interactions known to those of skill in the art. After contacting the recombinant cell for a sufficient time to effect any interactions, the level of gene expression is measured. The amount of time to effect such interactions may be empirically determined, such as by running a time course and measuring the level of transcription as a function of time. The amount of transcription may be measured using any method known to those of skill in the
25 art to be suitable. For example, specific mRNA expression may be detected using Northern blots or specific protein product may be identified by a characteristic stain. The amount of transcription is then compared to the amount of transcription in either the same cell in the absence of the test compound or it may be compared with the amount of transcription in a substantially identical cell that lacks the specific receptors. A substantially identical cell
30 may be derived from the same cells from which the recombinant cell was prepared but which had not been modified by introduction of heterologous DNA. Alternatively, it may be a cell in which the specific receptors are removed. Any statistically or otherwise

significant difference in the amount of transcription indicates that the test compound has in some manner altered the activity of the specific receptor.

If the test compound does not appear to enhance, activate or induce the activity of the cell surface protein, the assay may be repeated and modified by the introduction of a
5 step in which the recombinant cell is first tested for the ability of a known agonist or activator of the specific receptor to activate transcription if the transcription is induced, the test compound is then assayed for its ability to inhibit, block or otherwise affect the activity of the agonist.

The transcription based assay is useful for identifying compounds that interact with
10 any cell surface protein whose activity ultimately alters gene expression. In particular, the assays can be used to test functional ligand-receptor or ligand-ion channel interactions for a number of categories of cell surface-localized receptors, including: ligand-gated ion channels and voltage-gated ion channels, and G protein-coupled receptors.

Any transfectable cell that can express the desired cell surface protein in a manner
15 such the protein functions to intracellularly transduce an extracellular signal may be used. The cells may be selected such that they endogenously express the cell surface protein or may be genetically engineered to do so. Many such cells are known to those of skill in the art. Such cells include, but are not limited to Ltk< - > cells, PC12 cells and COS-7 cells.

The preparation of cells which express a receptor or ion channel and a reporter gene
20 expression construct, and which are useful for testing compounds to assess their activities, is exemplified in the Examples provided herewith by reference to mammalian Ltk< - > and COS-7 cell lines, which express the Type I human muscarinic (HM1) receptor and which are transformed with either a c-fos promoter-CAT reporter gene expression construct or a c-fos promoter-luciferase reporter gene expression construct.

25 Any cell surface protein that is known to those of skill in the art or that may be identified by those of skill in the art may used in the assay. The cell surface protein may endogenously expressed on the selected cell or it may be expressed from cloned DNA. Exemplary cell surface proteins include, but are not limited to, cell surface receptors and ion channels. Cell surface receptors include, but are not limited to, muscarinic receptors
30 (e.g., human M2 (GenBank accession #M16404); rat M3 (GenBank accession #M16407); human M4 (GenBank accession #M16405); human M5 (Bonner et al. (1988) Neuron 1:403-410); and the like); neuronal nicotinic acetylcholine receptors (e.g., the alpha 2, alpha 3 and beta 2 subtypes disclosed in U.S. Ser. No. 504,455 (filed Apr. 3, 1990), hereby

expressly incorporated by reference herein in its entirety); the rat alpha 2 subunit (Wada et al. (1988) Science 240:330-334); the rat alpha 3 subunit (Boulter et al. (1986) Nature 319:368-374); the rat alpha 4 subunit (Goldman et al. (1987) cell 48:965-973); the rat alpha 5 subunit (Boulter et al. (1990) J. Biol. Chem. 265:4472-4482); the rat beta 2 subunit
5 (Deneris et al. (1988) Neuron 1:45-54); the rat beta 3 subunit (Deneris et al. (1989) J. Biol. Chem. 264: 6268-6272); the rat beta 4 subunit (Duvoisin et al. (1989) Neuron 3:487-496); combinations of the rat alpha subunits, beta subunits and alpha and beta subunits; GABA receptors (e.g., the bovine alpha 1 and beta 1 subunits (Schofield et al. (1987) Nature 328:221-227); the bovine alpha 2 and alpha 3 subunits (Levitan et al. (1988) Nature 335:76-
10 79); the gamma -subunit (Pritchett et al. (1989) Nature 338:582-585); the beta 2 and beta 3 subunits (Ymer et al. (1989) EMBO J. 8:1665-1670); the delta subunit (Shivers, B.D. (1989) Neuron 3:327-337); and the like); glutamate receptors (e.g., receptor isolated from rat brain (Hollmann et al. (1989) Nature 342:643-648); and the like); adrenergic receptors (e.g., human beta 1 (Frielle et al. (1987) Proc. Natl. Acad. Sci. 84.:7920-7924); human
15 alpha 2 (Kobilka et al. (1987) Science 238:650-656); hamster beta 2 (Dixon et al. (1986) Nature 321:75-79); and the like); dopamine receptors (e.g., human D2 (Stormann et al. (1990) Molec. Pharm.37:1-6); rat (Bunzow et al. (1988) Nature 336:783-787); and the like); NGF receptors (e.g., human NGF receptors (Johnson et al. (1986) Cell 47:545-554); and the like); serotonin receptors (e.g., human 5HT1a (Kobilka et al. (1987) Nature 329:75-79); rat
20 5HT2 (Julius et al. (1990) PNAS 87:928-932); rat 5HT1c (Julius et al. (1988) Science 241:558-564); and the like).

Reporter gene constructs are prepared by operatively linking a reporter gene with at least one transcriptional regulatory element. If only one transcriptional regulatory element is included, it must be a regulatable promoter. At least one of the selected transcriptional
25 regulatory elements must be indirectly or directly regulated by the activity of the selected cell-surface receptor whereby activity of the receptor can be monitored via transcription of the reporter genes.

The construct may contain additional transcriptional regulatory elements, such as a FIRE sequence, or other sequence, that is not necessarily regulated by the cell surface
30 protein, but is selected for its ability to reduce background level transcription or to amplify the transduced signal and to thereby increase the sensitivity and reliability of the assay.

Many reporter genes and transcriptional regulatory elements are known to those of skill in the art and others may be identified or synthesized by methods known to those of skill in the art.

A reporter gene includes any gene that expresses a detectable gene product, which
5 may be RNA or protein. Preferred reporter genes are those that are readily detectable. The reporter gene may also be included in the construct in the form of a fusion gene with a gene that includes desired transcriptional regulatory sequences or exhibits other desirable properties.

Examples of reporter genes include, but are not limited to CAT (chloramphenicol
10 acetyl transferase) (Alton and Vapnek (1979), Nature 282: 864-869) luciferase, and other enzyme detection systems, such as beta-galactosidase; firefly luciferase (deWet et al. (1987), Mol. Cell. Biol. 7:725-737); bacterial luciferase (Engbrecht and Silverman (1984), PNAS 1: 4154-4158; Baldwin et al. (1984), Biochemistry 23: 3663-3667); alkaline phosphatase (Toh et al. (1989) Eur. J. Biochem. 182: 231-238, Hall et al. (1983) J. Mol.
15 Appl. Gen. 2: 101).

Transcriptional control elements include, but are not limited to, promoters, enhancers, and repressor and activator binding sites. Suitable transcriptional regulatory elements may be derived from the transcriptional regulatory regions of genes whose expression is rapidly induced, generally within minutes, of contact between the cell surface
20 protein and the effector protein that modulates the activity of the cell surface protein. Examples of such genes include, but are not limited to, the immediate early genes (see, Sheng et al. (1990) Neuron 4: 477-485), such as c-fos, Immediate early genes are genes that are rapidly induced upon binding of a ligand to a cell surface protein. The transcriptional control elements that are preferred for use in the gene constructs include transcriptional
25 control elements from immediate early genes, elements derived from other genes that exhibit some or all of the characteristics of the immediate early genes, or synthetic elements that are constructed such that genes in operative linkage therewith exhibit such characteristics. The characteristics of preferred genes from which the transcriptional control elements are derived include, but are not limited to, low or undetectable expression in
30 quiescent cells, rapid induction at the transcriptional level within minutes of extracellular stimulation, induction that is transient and independent of new protein synthesis, subsequent shut-off of transcription requires new protein synthesis, and mRNAs transcribed from these genes have a short half-life. It is not necessary for all of these properties to be present.

Pharmaceutical Compositions/Formulations

In another aspect, the present invention provides pharmaceutically acceptable compositions which comprise a therapeutically-effective amount of one or more of the compounds described above, formulated together with one or more pharmaceutically acceptable carriers (additives) and/or diluents. As described in detail below, the pharmaceutical compositions of the present invention may be specially formulated for administration in solid or liquid form, including those adapted for the following: (1) oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, e.g., those targeted for buccal, sublingual, and systemic absorption, boluses, powders, granules, pastes for application to the tongue, hard gelatin capsules, soft gelatin capsules, mouth sprays, syrups, emulsions, micro-emulsions; (2) parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; (3) topical application, for example, as a cream, ointment, or a controlled-release patch or spray applied to the skin; (4) intravaginally or intrarectally, for example, as a pessary, cream or foam; (5) sublingually; (6) ocularly; (7) transdermally; or (8) nasally.

The phrase "therapeutically-effective amount" as used herein means that amount of a compound, material, or composition comprising a compound of the present invention which is effective for producing some desired therapeutic effect in at least a sub-population of cells in an animal at a reasonable benefit/risk ratio applicable to any medical treatment.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The phrase "pharmaceutically-acceptable carrier" as used herein means a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, manufacturing aid (e.g., lubricant, talc magnesium, calcium or zinc stearate, or stearic acid), or solvent encapsulating material, involved in carrying or transporting the subject compound from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically-acceptable carriers include: (1)

sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) pH buffered solutions; (21) polyesters, polycarbonates and/or polyanhydrides; and (22) other non-toxic compatible substances employed in pharmaceutical formulations.

As set out above, certain embodiments of the present compounds may contain a basic functional group, such as amino or alkylamino, and are, thus, capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable acids. The term "pharmaceutically-acceptable salts" in this respect, refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared *in situ* in the administration vehicle or the dosage form manufacturing process, or by separately reacting a purified compound of the invention in its free base form with a suitable organic or inorganic acid, and isolating the salt thus formed during subsequent purification. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, besylate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts and the like. (See, for example, Berge et al. (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19)

The pharmaceutically acceptable salts of the subject compounds include the conventional nontoxic salts or quaternary ammonium salts of the compounds, e.g., from non-toxic organic or inorganic acids. For example, such conventional nontoxic salts include those derived from inorganic acids such as hydrochloride, hydrobromic, sulfuric, sulfamic, phosphoric, nitric, and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, palmitic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, benzenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isothionic, and the like.

In other cases, the compounds of the present invention may contain one or more acidic functional groups and, thus, are capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable bases. The term "pharmaceutically-acceptable salts" in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention. These salts can likewise be prepared *in situ* in the administration vehicle or the dosage form manufacturing process, or by separately reacting the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically-acceptable metal cation, with ammonia, or with a pharmaceutically-acceptable organic primary, secondary or tertiary amine. Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like. (See, for example, Berge et al., *supra*)

Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

Examples of pharmaceutically-acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

Formulations of the present invention include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will vary depending upon the host being treated, the particular mode of administration. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will generally be that amount of the compound which produces a therapeutic

effect. Generally, out of one hundred per cent, this amount will range from about 0.1 per cent to about ninety-nine percent of active ingredient, preferably from about 5 per cent to about 70 per cent, most preferably from about 10 per cent to about 30 per cent.

Methods of preparing these formulations or compositions include the step of
5 bringing into association a compound of the present invention with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a compound of the present invention with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

10 Formulations of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia)
15 and/or as mouth washes and the like, each containing a predetermined amount of a compound of the present invention as an active ingredient. A compound of the present invention may also be administered as a bolus, electuary or paste.

In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules, trouches and the like), the active ingredient is mixed with
20 one or more pharmaceutically-acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium
25 carbonate, potato or tapioca starch, alginic acid, corn starch, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds, and surfactants, such as poloxamer and sodium lauryl sulfate; (7) wetting agents, such as, for example, cetyl alcohol, glycerol monostearate, and non-ionic surfactants; (8) absorbents, such as kaolin and bentonite clay;
30 (9) lubricants, such a talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, zinc stearate, sodium stearate, stearic acid and mixtures thereof; (10) coloring agents; and (11) controlled release agents, such as crospovidone or ethyl cellulose. In the case of capsules, tablets and pills, the pharmaceutical compositions may also

comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-shelled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding, optionally with one or more
5 accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid
10 diluent.

The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or
15 controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be formulated for rapid release, e.g., freeze-dried. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which
20 can be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The
25 active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Liquid dosage forms for oral administration of the compounds of the invention include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain
30 inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils),

glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring,
5 perfuming and preservative agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

10 Formulations of the pharmaceutical compositions of the invention for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more compounds of the invention with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body
15 temperature and, therefore, will melt in the rectum or vaginal cavity and release the active compound.

Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, vaginal tablets, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

20 Dosage forms for the topical or transdermal administration of a compound of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically-acceptable carrier, and with any preservatives, buffers, or propellants which may be required.

25 The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

Powders and sprays can contain, in addition to a compound of this invention,
30 excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

Transdermal patches have the added advantage of providing controlled delivery of a compound of the present invention to the body. Such dosage forms can be made by dissolving or dispersing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux can
5 be controlled by either providing a rate controlling membrane or dispersing the compound in a polymer matrix or gel.

Ophthalmic formulations, eye ointments, powders, solutions and the like, are also contemplated as being within the scope of this invention.

Pharmaceutical compositions of this invention suitable for parenteral administration
10 comprise one or more compounds of the invention in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain sugars, alcohols, antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the
15 blood of the intended recipient or suspending or thickening agents.

Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper
20 fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of
25 microorganisms upon the subject compounds may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which
30 delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having

poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form.

Alternatively, delayed absorption of a parenterally-administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

5 Injectable depot forms are made by forming microencapsule matrices of the subject compounds in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by
10 entrapping the drug in liposomes or microemulsions which are compatible with body tissue.

When the compounds of the present invention are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1 to 99% (more preferably, 10 to 30%) of active ingredient in combination with a pharmaceutically acceptable carrier.

15 The preparations of the present invention may be given orally, parenterally, topically, or rectally. They are of course given in forms suitable for each administration route. For example, they are administered in tablets or capsule form, by injection, inhalation, eye lotion, ointment, suppository, etc. administration by injection, infusion or inhalation; topical by lotion or ointment; and rectal by suppositories. Oral administrations
20 are preferred.

 The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal,
25 transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

 The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the central nervous system, such
30 that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

 These compounds may be administered to humans and other animals for therapy by any suitable route of administration, including orally, nasally, as by, for example, a spray,

rectally, intravaginally, parenterally, intracisternally and topically, as by powders, ointments or drops, including buccally and sublingually.

Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical
5 compositions of the present invention, are formulated into pharmaceutically-acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition,
10 and mode of administration, without being toxic to the patient.

The selected dosage level will depend upon a variety of factors including the activity of the particular compound of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion or metabolism of the particular compound being employed, the rate and extent of
15 absorption, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

A physician or veterinarian having ordinary skill in the art can readily determine
20 and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

25 In general, a suitable daily dose of a compound of the invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Generally, oral, intravenous, intracerebroventricular and subcutaneous doses of the compounds of this invention for a patient, when used for the indicated effects, will range from about 0.0001 to
30 about 100 mg per kilogram of body weight per day.

If desired, the effective daily dose of the active compound may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals

throughout the day, optionally, in unit dosage forms. Preferred dosing is one administration per day.

While it is possible for a compound of the present invention to be administered alone, it is preferable to administer the compound as a pharmaceutical formulation

5 (composition).

In another aspect, the present invention provides pharmaceutically acceptable compositions which comprise a therapeutically-effective amount of one or more of the subject compounds, as described above, formulated together with one or more pharmaceutically acceptable carriers (additives) and/or diluents. As described in detail
10 below, the pharmaceutical compositions of the present invention may be specially formulated for administration in solid or liquid form, including those adapted for the following: (1) oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, boluses, powders, granules, pastes for application to the tongue; (2) parenteral administration, for example, by subcutaneous, intramuscular or
15 intravenous injection as, for example, a sterile solution or suspension; (3) topical application, for example, as a cream, ointment or spray applied to the skin, lungs, or mucous membranes; or (4) intravaginally or intrarectally, for example, as a pessary, cream or foam; (5) sublingually or buccally; (6) ocularly; (7) transdermally; or (8) nasally.

The compounds according to the invention may be formulated for administration in
20 any convenient way for use in human or veterinary medicine, by analogy with other pharmaceuticals.

The term "treatment" is intended to encompass also prophylaxis, therapy and cure.

The patient receiving this treatment is any animal in need, including primates, in particular humans, and other mammals such as equines, cattle, swine and sheep; and poultry
25 and pets in general.

The compound of the invention can be administered as such or in admixtures with pharmaceutically acceptable carriers and can also be administered in conjunction with antimicrobial agents such as penicillins, cephalosporins, aminoglycosides and glycopeptides. Conjunctive therapy, thus includes sequential, simultaneous and separate
30 administration of the active compound in a way that the therapeutical effects of the first administered one is not entirely disappeared when the subsequent is administered.

The addition of the active compound of the invention to animal feed is preferably accomplished by preparing an appropriate feed premix containing the active compound in an effective amount and incorporating the premix into the complete ration.

Alternatively, an intermediate concentrate or feed supplement containing the active
5 ingredient can be blended into the feed. The way in which such feed premixes and complete rations can be prepared and administered are described in reference books (such as "Applied Animal Nutrition", W.H. Freedman and CO., San Francisco, U.S.A., 1969 or "Livestock Feeds and Feeding" O and B books, Corvallis, Ore., U.S.A., 1977).

Combinatorial Libraries

10 The subject reactions readily lend themselves to the creation of combinatorial libraries of compounds for the screening of pharmaceutical, agrochemical or other biological or medically-related activity or material-related qualities. A combinatorial library for the purposes of the present invention is a mixture of chemically related compounds which may be screened together for a desired property; said libraries may be in
15 solution or covalently linked to a solid support. The preparation of many related compounds in a single reaction greatly reduces and simplifies the number of screening processes which need to be carried out. Screening for the appropriate biological, pharmaceutical, agrochemical or physical property may be done by conventional methods.

Diversity in a library can be created at a variety of different levels. For instance, the
20 substrate aryl groups used in a combinatorial approach can be diverse in terms of the core aryl moiety, e.g., a variegation in terms of the ring structure, and/or can be varied with respect to the other substituents.

A variety of techniques are available in the art for generating combinatorial libraries of small organic molecules. See, for example, Blondelle et al. (1995) Trends Anal. Chem.
25 14:83; the Affymax U.S. Patents 5,359,115 and 5,362,899; the Ellman U.S. Patent 5,288,514; the Still et al. PCT publication WO 94/08051; Chen et al. (1994) JACS 116:2661; Kerr et al. (1993) JACS 115:252; PCT publications WO92/10092, WO93/09668 and WO91/07087; and the Lerner et al. PCT publication WO93/20242). Accordingly, a variety of libraries on the order of about 16 to 1,000,000 or more diversomers can be
30 synthesized and screened for a particular activity or property.

In an exemplary embodiment, a library of substituted diversomers can be synthesized using the subject reactions adapted to the techniques described in the Still et al. PCT publication WO 94/08051, e.g., being linked to a polymer bead by a hydrolyzable or

photolyzable group, e.g., located at one of the positions of substrate. According to the Still et al. technique, the library is synthesized on a set of beads, each bead including a set of tags identifying the particular diversomer on that bead. In one embodiment, which is particularly suitable for discovering enzyme inhibitors, the beads can be dispersed on the surface of a permeable membrane, and the diversomers released from the beads by lysis of the bead linker. The diversomer from each bead will diffuse across the membrane to an assay zone, where it will interact with an enzyme assay. Detailed descriptions of a number of combinatorial methodologies are provided below.

A. Direct Characterization

A growing trend in the field of combinatorial chemistry is to exploit the sensitivity of techniques such as mass spectrometry (MS), e.g., which can be used to characterize sub-femtomolar amounts of a compound, and to directly determine the chemical constitution of a compound selected from a combinatorial library. For instance, where the library is provided on an insoluble support matrix, discrete populations of compounds can be first released from the support and characterized by MS. In other embodiments, as part of the MS sample preparation technique, such MS techniques as MALDI can be used to release a compound from the matrix, particularly where a labile bond is used originally to tether the compound to the matrix. For instance, a bead selected from a library can be irradiated in a MALDI step in order to release the diversomer from the matrix, and ionize the diversomer for MS analysis.

B) Multipin Synthesis

The libraries of the subject method can take the multipin library format. Briefly, Geysen and co-workers (Geysen et al. (1984) PNAS 81:3998-4002) introduced a method for generating compound libraries by a parallel synthesis on polyacrylic acid-grated polyethylene pins arrayed in the microtitre plate format. The Geysen technique can be used to synthesize and screen thousands of compounds per week using the multipin method, and the tethered compounds may be reused in many assays. Appropriate linker moieties can also been appended to the pins so that the compounds may be cleaved from the supports after synthesis for assessment of purity and further evaluation (c.f., Bray et al. (1990) Tetrahedron Lett 31:5811-5814; Valerio et al. (1991) Anal Biochem 197:168-177; Bray et al. (1991) Tetrahedron Lett 32:6163-6166).

C) Divide-Couple-Recombine

In yet another embodiment, a variegated library of compounds can be provided on a set of beads utilizing the strategy of divide-couple-recombine (see, e.g., Houghten (1985) PNAS 82:5131-5135; and U.S. Patents 4,631,211; 5,440,016; 5,480,971). Briefly, as the name implies, at each synthesis step where degeneracy is introduced into the library, the beads are divided into separate groups equal to the number of different substituents to be added at a particular position in the library, the different substituents coupled in separate reactions, and the beads recombined into one pool for the next iteration.

In one embodiment, the divide-couple-recombine strategy can be carried out using an analogous approach to the so-called "tea bag" method first developed by Houghten, where compound synthesis occurs on resin sealed inside porous polypropylene bags (Houghten et al. (1986) PNAS 82:5131-5135). Substituents are coupled to the compound-bearing resins by placing the bags in appropriate reaction solutions, while all common steps such as resin washing and deprotection are performed simultaneously in one reaction vessel. At the end of the synthesis, each bag contains a single compound.

D) Combinatorial Libraries by Light-Directed, Spatially Addressable Parallel Chemical Synthesis

A scheme of combinatorial synthesis in which the identity of a compound is given by its locations on a synthesis substrate is termed a spatially-addressable synthesis. In one embodiment, the combinatorial process is carried out by controlling the addition of a chemical reagent to specific locations on a solid support (Dower et al. (1991) Annu Rep Med Chem 26:271-280; Fodor, S.P.A. (1991) Science 251:767; Pirrung et al. (1992) U.S. Patent No. 5,143,854; Jacobs et al. (1994) Trends Biotechnol 12:19-26). The spatial resolution of photolithography affords miniaturization. This technique can be carried out through the use protection/deprotection reactions with photolabile protecting groups.

The key points of this technology are illustrated in Gallop et al. (1994) J Med Chem 37:1233-1251. A synthesis substrate is prepared for coupling through the covalent attachment of photolabile nitroveratryloxycarbonyl (NVOC) protected amino linkers or other photolabile linkers. Light is used to selectively activate a specified region of the synthesis support for coupling. Removal of the photolabile protecting groups by light (deprotection) results in activation of selected areas. After activation, the first of a set of amino acid analogs, each bearing a photolabile protecting group on the amino terminus, is exposed to the entire surface. Coupling only occurs in regions that were addressed by light in the preceding step. The reaction is stopped, the plates washed, and the substrate is again

illuminated through a second mask, activating a different region for reaction with a second protected building block. The pattern of masks and the sequence of reactants define the products and their locations. Since this process utilizes photolithography techniques, the number of compounds that can be synthesized is limited only by the number of synthesis sites that can be addressed with appropriate resolution. The position of each compound is precisely known; hence, its interactions with other molecules can be directly assessed.

In a light-directed chemical synthesis, the products depend on the pattern of illumination and on the order of addition of reactants. By varying the lithographic patterns, many different sets of test compounds can be synthesized simultaneously; this characteristic leads to the generation of many different masking strategies.

E) Encoded Combinatorial Libraries

In yet another embodiment, the subject method utilizes a compound library provided with an encoded tagging system. A recent improvement in the identification of active compounds from combinatorial libraries employs chemical indexing systems using tags that uniquely encode the reaction steps a given bead has undergone and, by inference, the structure it carries. Conceptually, this approach mimics phage display libraries, where activity derives from expressed peptides, but the structures of the active peptides are deduced from the corresponding genomic DNA sequence. The first encoding of synthetic combinatorial libraries employed DNA as the code. A variety of other forms of encoding have been reported, including encoding with sequenceable bio-oligomers (e.g., oligonucleotides and peptides), and binary encoding with additional non-sequenceable tags.

1) Tagging with sequenceable bio-oligomers

The principle of using oligonucleotides to encode combinatorial synthetic libraries was described in 1992 (Brenner et al. (1992) PNAS 89:5381-5383), and an example of such a library appeared the following year (Needles et al. (1993) PNAS 90:10700-10704). A combinatorial library of nominally 7^7 (= 823,543) peptides composed of all combinations of Arg, Gln, Phe, Lys, Val, D-Val and Thr (three-letter amino acid code), each of which was encoded by a specific dinucleotide (TA, TC, CT, AT, TT, CA and AC, respectively), was prepared by a series of alternating rounds of peptide and oligonucleotide synthesis on solid support. In this work, the amine linking functionality on the bead was specifically differentiated toward peptide or oligonucleotide synthesis by simultaneously preincubating the beads with reagents that generate protected OH groups for oligonucleotide synthesis and protected NH₂ groups for peptide synthesis (here, in a ratio

of 1:20). When complete, the tags each consisted of 69-mers, 14 units of which carried the code. The bead-bound library was incubated with a fluorescently labeled antibody, and beads containing bound antibody that fluoresced strongly were harvested by fluorescence-activated cell sorting (FACS). The DNA tags were amplified by PCR and sequenced, and the predicted peptides were synthesized. Following such techniques, compound libraries can be derived for use in the subject method, where the oligonucleotide sequence of the tag identifies the sequential combinatorial reactions that a particular bead underwent, and therefore provides the identity of the compound on the bead.

The use of oligonucleotide tags permits exquisitely sensitive tag analysis. Even so, the method requires careful choice of orthogonal sets of protecting groups required for alternating co-synthesis of the tag and the library member. Furthermore, the chemical lability of the tag, particularly the phosphate and sugar anomeric linkages, may limit the choice of reagents and conditions that can be employed for the synthesis of non-oligomeric libraries. In preferred embodiments, the libraries employ linkers permitting selective detachment of the test compound library member for assay.

Peptides have also been employed as tagging molecules for combinatorial libraries. Two exemplary approaches are described in the art, both of which employ branched linkers to solid phase upon which coding and ligand strands are alternately elaborated. In the first approach (Kerr JM et al. (1993) J Am Chem Soc 115:2529-2531), orthogonality in synthesis is achieved by employing acid-labile protection for the coding strand and base-labile protection for the compound strand.

In an alternative approach (Nikolaiev et al. (1993) Pept Res 6:161-170), branched linkers are employed so that the coding unit and the test compound can both be attached to the same functional group on the resin. In one embodiment, a cleavable linker can be placed between the branch point and the bead so that cleavage releases a molecule containing both code and the compound (Ptek et al. (1991) Tetrahedron Lett 32:3891-3894). In another embodiment, the cleavable linker can be placed so that the test compound can be selectively separated from the bead, leaving the code behind. This last construct is particularly valuable because it permits screening of the test compound without potential interference of the coding groups. Examples in the art of independent cleavage and sequencing of peptide library members and their corresponding tags has confirmed that the tags can accurately predict the peptide structure.

2) Non-sequenceable Tagging: Binary Encoding

An alternative form of encoding the test compound library employs a set of non-sequencable electrophoric tagging molecules that are used as a binary code (Ohlmeyer et al. (1993) PNAS 90:10922-10926). Exemplary tags are haloaromatic alkyl ethers that are detectable as their trimethylsilyl ethers at less than femtomolar levels by electron capture gas chromatography (ECGC). Variations in the length of the alkyl chain, as well as the nature and position of the aromatic halide substituents, permit the synthesis of at least 40 such tags, which in principle can encode 2^{40} (e.g., upwards of 10^{12}) different molecules. In the original report (Ohlmeyer et al., supra) the tags were bound to about 1% of the available amine groups of a peptide library via a photocleavable α -nitrobenzyl linker. This approach is convenient when preparing combinatorial libraries of peptide-like or other amine-containing molecules. A more versatile system has, however, been developed that permits encoding of essentially any combinatorial library. Here, the compound would be attached to the solid support via the photocleavable linker and the tag is attached through a catechol ether linker via carbene insertion into the bead matrix (Nestler et al. (1994) J Org Chem 59:4723-4724). This orthogonal attachment strategy permits the selective detachment of library members for assay in solution and subsequent decoding by ECGC after oxidative detachment of the tag sets.

Although several amide-linked libraries in the art employ binary encoding with the electrophoric tags attached to amine groups, attaching these tags directly to the bead matrix provides far greater versatility in the structures that can be prepared in encoded combinatorial libraries. Attached in this way, the tags and their linker are nearly as unreactive as the bead matrix itself. Two binary-encoded combinatorial libraries have been reported where the electrophoric tags are attached directly to the solid phase (Ohlmeyer et al. (1995) PNAS 92:6027-6031) and provide guidance for generating the subject compound library. Both libraries were constructed using an orthogonal attachment strategy in which the library member was linked to the solid support by a photolabile linker and the tags were attached through a linker cleavable only by vigorous oxidation. Because the library members can be repetitively partially photoeluted from the solid support, library members can be utilized in multiple assays. Successive photoelution also permits a very high throughput iterative screening strategy: first, multiple beads are placed in 96-well microtiter plates; second, compounds are partially detached and transferred to assay plates; third, a metal binding assay identifies the active wells; fourth, the corresponding beads are

rearrayed singly into new microtiter plates; fifth, single active compounds are identified; and sixth, the structures are decoded.


Exemplification

The invention now being generally described, it will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

Example 1

SYNTHESIS OF 1-[1-(4-CHLORO-PHENYL)-CYCLOBUTYL]-(4-HYDROXY-PIPERIDIN-1-YL)-

METHADONE




To a stirred solution of piperidin-4-ol (5.0 g, 0.049 moles) and 1-(4-chloro-phenyl)-cyclobutane carboxylic acid (15.60 g, 0.074 moles) in anhydrous DCM (100 mL) was added di-isopropyl ethyl amine (25.8 mL, 0.148 moles) dropwise. After completion of addition solid PyBroP (34.3 g, 0.074 moles) was added to the stirring reaction mixture. The reaction mixture continued stirring at RT for 10h and was quenched with 10% KOH (aq.). The aqueous layer was extracted with EtOAc 3x (200 mL). Combined organic layers were dried over Na₂SO₄ and concentrated to yield an oil. This crude material was purified using silica gel chromatography (1:1 hexane:EtOAc) to yield 1 as an oil (13 g, 0.044 mole, 90%).

LRMS: M+ 294.

Example 2

Synthesis of 1-[1-(4-Chloro-phenyl)-cyclobutylmethyl]-piperidin-4-ol




A solution of amide 1 (1 g, 3.40 mmoles) dissolved in anhydrous toluene (17 mL) was cooled to 0° C. RedAl (2.41 g, 11.91 mmoles) was added dropwise to the cooled stirring reaction mixture. After completion of addition the reaction continued stirring at RT.

- 5 After 1.5h water was added to the reaction mixture. The aqueous layer was extracted with EtOAc 3x (50 mL). Combined organic layers were dried over Na₂SO₄ and concentrated to yield 2 an oil. LRMS: M+ 282.


Example 3

- 10 Synthesis of [4-(3,5-bis-trifluoromethyl-phenoxy)piperidin-1-yl]-[1-(4-chloro-phenyl)-cyclobutyl]-methanone


To a stirring solution of 1 (0.50 g, 1.70 mmole) in anhydrous DMF



- 50 mL) was added solid NaH (122 mg, 5.10 mmole). After completion of addition the reaction mixture stirred for 45 min. at RT before the 3,5-bis(trifluoromethyl) benzyl bromide (583 mg, 1.9 mmole) was added. The reaction progress was monitored by HPLC and once complete was diluted with EtOAc and brine (5 mL). The aqueous layer was extracted with EtOAc (3 x 10 mL) and combined organics were dried over Na₂SO₄ and concentrated to yield an oil. The crude material was purified using silica gel chromatography (100% hexanes-80:20: hexanes: EtOAc) to yield 3 as an oil. LRMS: M+521.
- 15

Example 4**Synthesis of [1-(4-chloro-phenyl)-cyclobutyl]-[4-(4-trifluoromethyl-benzyloxy)piperidin-1-yl]-methanone**

5 To a stirring solution of 1 (1.0 g, 3.40 mmole) in anhydrous DMF (17 mL) was added solid NaH (245 mg, 10.2 mmole). After completion of addition the reaction mixture stirred for 45 min. at RT before the 4-trifluoromethyl benzyl bromide (884 mg, 3.7 mmole) was added. The reaction progress was monitored by HPLC and once complete was diluted with EtOAc and brine (5 mL) was added. The aqueous layer was extracted with EtOAc (3x,
10 20 mL) and combined organics were dried over Na₂SO₄ and concentrated to yield an oil. The crude material was purified using silica gel chromatography (100% hexanes-60:40: hexanes: EtOAc) to yield 4 as an oil. LRMS: M+451.


Example 5**Synthesis of [4-(3,5-bis-trifluoromethyl-benzyloxy)-1-[1-(4-chloro-phenyl)-cyclobutylmethyl]-piperidine**

A solution of amide 4 (560 mg, 1.08 mmoles) dissolved in anhydrous toluene (5.4 mL) was cooled to 0° C. RedAl (762 mg, 3.77 mmoles) was added dropwise to the cooled stirring reaction mixture. After completion of addition the reaction continued stirring at RT.
20 After 1.5h water was added to the reaction mixture. The aqueous layer was extracted with

EtOAc (3 x 10 mL). Combined organic layers were dried over Na₂SO₄ and concentrated to yield an oil. The crude material was purified using silica gel chromatography (99:1 CH₂Cl₂: 2M NH₃ in EtOH) to yield 5. LRMS: M+ 505.

Example 6


5 **Synthesis of [1-(4-chloro-phenyl)-cyclobutylmethyl]-4-(4-trifluoromethyl-benzyloxy)-piperidine**



A solution of amide 3 (350 mg, 0.77 mmoles) dissolved in anhydrous toluene (3.87 mL) was cooled to 0° C. RedAl (548 mg, 2.71 mmoles) was added dropwise to the cooled stirring reaction mixture. After completion of addition the reaction continued stirring at RT. After 1.5h water was added to the reaction mixture. The aqueous layer was extracted with EtOAc 3x (10 mL). Combined organic layers were dried over Na₂SO₄ and concentrated to yield an oil. The crude material was purified using silica gel chromatography (96:4 CH₂Cl₂: 2M NH₃ in EtOH) to yield 6. LRMS: M+ 439.

15 **Example 7**


Synthesis of [1-(4-chloro-phenyl)-cyclobutyl]-[4-(4-trifluoromethyl-phenoxy)piperidin-1-yl]-methanone



A solution of 1 (1.5 g, 5.10 mmoles), triphenylphosphine (2.0 g, 7.50 mmole), and phenol (1.65 g, 10.20 mmoles) dissolved in anhydrous ether (15.5 mL) was cooled in a brine bath to -5°C . DEAD (1.33 g, 7.60 mmoles) dissolved in ether (10 mL) was added to the cooled stirring reaction mixture. After completion of addition the reaction mixture continued stirring at -5°C . After 4h the reaction mixture was concentrated and crude material was dissolved in a hexane/ethyl acetate mixture (70% hexanes:30% ethyl acetate, 30 mL). Phosphine by-products precipitated and were filtered off. The filtrate was concentrated to yield an oil. This oil was purified using silica gel chromatography (100% hexanes-100% EtOAc) to yield the amine 7 (410 mg, 0.932 mmole, 18%). LRMS: $M+ 438$.

10 Example 8

Synthesis of 1-[1-(4-chloro-phenyl)-cyclobutylmethyl]-4-(4-trifluoromethyl-phenoxy)-piperidine




A solution of amide 7 (100 mg, 0.29 mmoles) dissolved in anhydrous toluene (1.14 mL) was cooled to 0°C . RedAl (161 mg, 0.80 mmoles) was added dropwise to the cooled stirring reaction mixture. After completion of addition the reaction continued stirring at RT. After 1.5h water was added to the reaction mixture. The aqueous layer was extracted with EtOAc (3 x 3 mL). Combined organic layers were dried over Na_2SO_4 and concentrated to yield an oil. The crude material was purified using silica gel chromatography (3:2

20 Hexanes:Et₂O) to yield 8. LRMS: $M+ 423$.

Example 9

Synthesis of 1-[1-(4-chloro-phenyl)-cyclobutanecarbonyl]-piperidine-4-carboxylic acid ethyl ester




To a stirred solution of piperidine-4-carboxylic acid ethyl ester (5.0 g, 0.032 moles) and 1-(4-chloro-phenyl)-cyclobutane carboxylic acid (9.98 g, 0.048 moles) in anhydrous DCM (100 mL) was added di-isopropyl ethyl amine (16.6 mL, 0.095 moles) dropwise.

- 5 After completion of addition solid PyBroP (22.6 g, 0.048 moles) was added to the stirring reaction mixture. The reaction mixture continued stirring at RT for 10h and was quenched with water. The aqueous layer was extracted with EtOAc 3x (200 mL). Combined organic layers were dried over Na₂SO₄ and concentrated to yield an oil. This crude material was purified using silica gel chromatography (100% hexanes-6:4 hexane:EtOAc) to yield 9 as
- 10 an oil (7.21 g, 65%).

Example 10

Synthesis of {1-[1-(4-chloro-phenyl)-cyclobuylmethyl]-piperidin-4-yl}-methanol




- To a cooled (0° C) solution of LAH (17.2 mL, 17.2 mmol, 1M solution in THF) in anhydrous THF was added ester 9 (2 g, 5.73 mmole) dissolved in anhydrous THF (28 mL).
- 15 After completion of addition the reaction continued stirring at RT. After 4 h the reaction mixture was quenched with 10% HCl (aq.) and the crude reaction mixture was basified using 10% NaOH. The aqueous layer was extracted with EtOAc (3x, 200 mL) and combined organic layers were dried over Na₂SO₄ and then concentrated to yield an oil. The

crude material was purified using silica gel chromatography (100% EtOAc) to yield **10** (1.3 g, 77%). LRMS: M+ 294.

Example 11


Synthesis of 1-[1-(4-chloro-phenyl)-cyclobutylmethyl]-4-(4-trifluoromethyl-phenoxy)-piperidine

5




To a stirring solution of **10** (0.20 g, 0.680 mmole) in anhydrous DMF (3.4 mL) was added solid NaH (49 mg, 2.04 mmole). After completion of addition the reaction mixture stirred for 45 min. at RT before the 4-trifluoromethyl benzyl bromide (179 mg, 0.748 mmole) was added. The reaction progress was monitored by HPLC and once complete was diluted with EtOAc and brine (2 mL) was added. The aqueous layer was extracted with EtOAc (3x, 8 mL) and combined organics were dried over Na₂SO₄ and concentrated to yield an oil. The crude material was purified using silica gel chromatography (95:5 hexanes: Et₂O w/ 1% 2M NH₃ in EtOH) to yield **11** as an oil. LRMS: M+452.

10

Example 12**SYNTHESIS OF METHANESULFONIC ACID 1-[1-(4-CHLORO-PHENYL)-CYCLOBUTYLMETHYL]-PIPERIDIN-4-YLMETHYL ESTER**

- 5 To a solution of primary alcohol 10 (800 mg, 2.72 mmol) in DCM (13 mL) at room temperature was added $i\text{Pr}_2\text{NEt}$ (1.2 mL, 6.8 mmol) followed by MsCl (0.25 mL, 3.0 mmol). The reaction mixture was allowed to stir for one hour before purifying the crude mixture using a presaturated silica gel plug (4:1 hexanes: EtOAc) to provide the desired mesylate 12 (330 mg, 33%). LRMS: M^+ 373.

Example 13**SYNTHESIS OF 1-[1-(4-CHLORO-PHENYL)-CYCLOBUTYLMETHYL]-4-(4-TRIFLUOROMETHYL-PHENOXYMETHYL)-PIPERIDINE**

- 15 To the mesylate 12 (330 mg, 0.89 mmol) in anhydrous acetonitrile (4.4 mL) was added α, α, α -trifluoro- p -cresol (144 mg, 0.89 mmol) followed by Cs_2CO_3 (1.45 g, 4.44 mmol). The reaction mixture was heated to 75 °C and the reaction progress was monitored

by HPLC. Upon completion the reaction mixture was quenched with water and the aqueous layer was extracted with EtOAc (3 x 20 mL). The Combined organic layer was then dried over Na₂SO₄, filtered and concentrated *in vacuo*. The resulting residue was purified by silica gel chromatography using a gradient (100% hexanes-100% EtOAc) to obtain the desired ether 13. LRMS: M+ 438.

Example 14

Antagonism of Monoamine Transporters

The ability of compounds of the present invention to displace norepinephrine ligands *in vitro* was determined by the methods of Galli *et al* (*J. Exp.Biol.* 198:2197, 1995) using desipramine (IC₅₀=920 nM) as a reference compound. The displacement of dopamine and serotonin ligands *in vitro* was determined by the methods of Gu *et al* (*J. Biol. Chem.* 269:7124, 1994) using GBR-12909 (IC₅₀(NE uptake) = 65 nM; IC₅₀(5-HT uptake) = 110 nM) as a reference compound.

<u>Compound</u>	<u>Uptake Profile (% inhibition @ 1 μM)</u>		
	NE uptake	DA uptake	5-HT uptake
2	65	88	36
5	85	61	14
6	78	85	9
8	38	25	56
10	62	70	21
11	97	97	87
13	97	94	83

Incorporation by Reference


All of the patents and publications cited herein are hereby incorporated by reference.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

We claim:

1. A compound represented by A:



A

wherein

- 5 X represents O, S, SO, SO₂, NR₇, NC(O)R₇, NC(O)OR₇, or NS(O)₂R₇;
 m is 0, 1, 2, 3, or 4;
 y is 0, 1, or 2;
 z is 0, 1, or 2;
 R₄ represents H, alkyl, cycloalkyl, alkenyl, aryl, or heteroaryl;
 10 R₅ and R₆ are selected independently for each occurrence from the group consisting
 of H, alkyl, aryl, heteroaryl, F, OR₁₁, and OC(O)R₁₁; or an instance of CR₅R₆ taken together
 is C(O);
 R₇ represents alkyl, cycloalkyl, aryl, heteroaryl, aralkyl, or heteroaralkyl;
 R₈ and R₉ are selected independently for each occurrence from the group consisting
 15 of H, alkyl, aryl, heteroaryl, F, OR₁₁, and OC(O)R₁₁; or an instance of CR₈R₉ taken together
 is C(O);
 R₁₀ represents H, alkyl, cycloalkyl, alkenyl, aryl, or heteroaryl;
 R₁₁ represents H, alkyl, cycloalkyl, alkenyl, aryl, or heteroaryl;
 a covalent bond may connect R₄ and an instance of R₅ or R₆;
 20 a covalent bond may connect R₁₀ and an instance of R₈ or R₉;
 any two instances of R₅ and R₆ may be connected through a covalent bond;

any two geminal or vicinal instances of R₈ and R₉ may be connected through a covalent bond; and

the stereochemical configuration at any stereocenter of a compound represented by A is *R*, *S*, or a mixture of these configurations.

- 5 2. The compound of claim 1, wherein X is O.
3. The compound of claim 1, wherein m is 2.
4. The compound of claim 1, wherein y is 0.
5. The compound of claim 1, wherein y is 1.
6. The compound of claim 1, wherein z is 0.
- 10 7. The compound of claim 1, wherein z is 1.
8. The compound of claim 1, wherein R₄ is aryl.
9. The compound of claim 1, wherein R₄ is 4-chlorophenyl.
10. The compound of claim 1, wherein R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl.
- 15 11. The compound of claim 1, wherein R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl.
12. The compound of claim 1, wherein R₁₀ is aryl.
13. The compound of claim 1, wherein R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl.
- 20 14. The compound of claim 1, wherein X is O; and R₄ is aryl.
15. The compound of claim 1, wherein X is O; R₄ is aryl; and R₁₀ is aryl.
16. The compound of claim 1, wherein X is O; R₄ is aryl; R₁₀ is aryl; and R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl.
17. The compound of claim 1, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are
25 selected independently for each occurrence from the group consisting of H, and alkyl; and
 R₈ and R₉ are selected independently for each occurrence from the group consisting of H,
 and alkyl.
18. The compound of claim 1, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are
 selected independently for each occurrence from the group consisting of H, and alkyl; R₈
30 and R₉ are selected independently for each occurrence from the group consisting of H, and
 alkyl; and m is 2.
19. The compound of claim 1, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are
 selected independently for each occurrence from the group consisting of H, and alkyl; R₈

and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; and y is 0.

20. The compound of claim 1, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; and y is 1.

21. The compound of claim 1, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; and z is 0.

22. The compound of claim 1, wherein X is O; R₄ is aryl; R₁₀ is aryl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; and z is 1.

23. The compound of claim 1, wherein X is O; R₄ is 4-chlorophenyl; and R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl.

24. The compound of claim 1, wherein X is O; R₄ is 4-chlorophenyl; R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; and R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl.

25. The compound of claim 1, wherein X is O; R₄ is 4-chlorophenyl; R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; and m is 2.

26. The compound of claim 1, wherein X is O; R₄ is 4-chlorophenyl; R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; y is 1; and z is 1.

27. The compound of claim 1, wherein X is O; R₄ is 4-chlorophenyl; R₁₀ is 4-(trifluoromethyl)phenyl or 3,5-bis(trifluoromethyl)phenyl; R₅ and R₆ are selected independently for each occurrence from the group consisting of H, and alkyl; R₈ and R₉ are

selected independently for each occurrence from the group consisting of H, and alkyl; m is 2; y is 1; and z is 0.

28. The compound of claim 1, wherein said compound has an EC₅₀ less than 1 μ M in an assay based on a mammalian dopamine, serotonin, or norepinephrine receptor or
5 transporter.
29. The compound of claim 1, wherein said compound has an EC₅₀ less than 100 nM in an assay based on a mammalian dopamine, serotonin, or norepinephrine receptor or transporter.
30. The compound of claim 1, wherein said compound has an EC₅₀ less than 10 nM in
10 an assay based on a mammalian dopamine, serotonin, or norepinephrine receptor or transporter.
31. The compound of claim 1, wherein said compound has an EC₅₀ less than 1 μ M in an assay based on a mammalian dopamine receptor or transporter.
32. The compound of claim 1, wherein said compound has an EC₅₀ less than 100 nM in
15 an assay based on a mammalian dopamine receptor or transporter.
33. The compound of claim 1, wherein said compound has an EC₅₀ less than 10 nM in an assay based on a mammalian dopamine receptor or transporter.
34. The compound of claim 1, wherein said compound has an IC₅₀ less than 1 μ M in an assay based on a mammalian dopamine, serotonin, or norepinephrine receptor or
20 transporter.
35. The compound of claim 1, wherein said compound has an IC₅₀ less than 100 nM in an assay based on a mammalian dopamine, serotonin, or norepinephrine receptor or transporter.
36. The compound of claim 1, wherein said compound has an IC₅₀ less than 10 nM in
25 an assay based on a mammalian dopamine, serotonin, or norepinephrine receptor or transporter.
37. The compound of claim 1, wherein said compound has an IC₅₀ less than 1 μ M in an assay based on a mammalian dopamine receptor or transporter.
38. The compound of claim 1, wherein said compound has an IC₅₀ less than 100 nM in
30 an assay based on a mammalian dopamine receptor or transporter.
39. The compound of claim 1, wherein said compound has an IC₅₀ less than 10 nM in an assay based on a mammalian dopamine receptor or transporter.
40. The compound of claim 1, wherein said compound is a single stereoisomer.

41. A formulation, comprising a compound of claim 1; and a pharmaceutically acceptable excipient.
42. A method of modulating the activity of a dopamine, serotonin, or norepinephrine receptor or transporter in a mammal, comprising the step of:
- 5 administering to said mammal a therapeutically effective amount of a compound of claim 1.
43. The method of claim 42, wherein said mammal is a primate, equine, canine or feline.
44. The method of claim 42, wherein said mammal is a human.
- 10 45. The method of claim 42, wherein said compound is administered orally.
46. The method of claim 42, wherein said compound is administered intravenously.
47. The method of claim 42, wherein said compound is administered sublingually.
48. The method of claim 42, wherein said compound is administered ocularly.
49. The method of claim 42, wherein said compound is administered transdermally.
- 15 50. The method of claim 42, wherein said compound is administered rectally.
51. The method of claim 42, wherein said compound is administered vaginally.
52. The method of claim 42, wherein said compound is administered topically.
53. The method of claim 42, wherein said compound is administered intramuscularly.
54. The method of claim 42, wherein said compound is administered subcutaneously.
- 20 55. The method of claim 42, wherein said compound is administered buccally.
56. The method of claim 42, wherein said compound is administered nasally.
57. A method of modulating the activity of a dopamine receptor or transporter in a mammal, comprising the step of:
- administering to said mammal a therapeutically effective amount of a
- 25 compound of claim 1.
58. The method of claim 57, wherein said mammal is a primate, equine, canine or feline.
59. The method of claim 57, wherein said mammal is a human.
60. The method of claim 57, wherein said compound is administered orally.
- 30 61. The method of claim 57, wherein said compound is administered intravenously.
62. The method of claim 57, wherein said compound is administered sublingually.
63. The method of claim 57, wherein said compound is administered ocularly.
64. The method of claim 57, wherein said compound is administered transdermally.

65. The method of claim 57, wherein said compound is administered rectally.
66. The method of claim 57, wherein said compound is administered vaginally.
67. The method of claim 57, wherein said compound is administered topically.
68. The method of claim 57, wherein said compound is administered intramuscularly.
- 5 69. The method of claim 57, wherein said compound is administered subcutaneously.
70. The method of claim 57, wherein said compound is administered buccally.
71. The method of claim 57, wherein said compound is administered nasally.
72. A method of treating a mammal suffering from addiction, anxiety, depression, sexual dysfunction, hypertension, migraine, Alzheimer's disease, obesity, emesis, 10 psychosis, analgesia, schizophrenia, Parkinson's disease, restless leg syndrome, sleeping disorders, attention deficit hyperactivity disorder, irritable bowel syndrome, premature ejaculation, menstrual dysphoria syndrome, urinary incontinence, inflammatory pain, neuropathic pain, Lesche-Nyhan disease, Wilson's disease, or Tourette's syndrome, comprising the step of:
- 15 administering to said mammal a therapeutically effective amount of a compound of claim 1.
73. The method of claim 72, wherein said mammal is a primate, equine, canine or feline.
74. The method of claim 72, wherein said mammal is a human.
- 20 75. The method of claim 72, wherein said compound is administered orally.
76. The method of claim 72, wherein said compound is administered intravenously.
77. The method of claim 72, wherein said compound is administered sublingually.
78. The method of claim 72, wherein said compound is administered ocularly.
79. The method of claim 72, wherein said compound is administered transdermally.
- 25 80. The method of claim 72, wherein said compound is administered rectally.
81. The method of claim 72, wherein said compound is administered vaginally.
82. The method of claim 72, wherein said compound is administered topically.
83. The method of claim 72, wherein said compound is administered intramuscularly.
84. The method of claim 72, wherein said compound is administered subcutaneously.
- 30 85. The method of claim 72, wherein said compound is administered buccally.
86. The method of claim 72, wherein said compound is administered nasally.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US02/39651

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : Please See Extra Sheet. US CL : 514/317, 326; 546/193, 207, 232 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/317, 326; 546/193, 207, 232 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS--structure EAST/WEST--subclass		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database CAS on STN (Columbus, OH, USA) , DN 133:105053, YAMADA et al. 'Preparation of acylpiperazines, acylpiperidines and related compounds as cholinergic agonists' WO 2000042011 (2000), see RN 283167-13-5.	1
X ----- Y	Database CAS on STN (Columbus, OH, USA), DN 126:47116, RIEMER, c. '[re[aratopm pf 1-benzylpiperidiny]benzoates and related compounds as antipsychotics' WO 9635666 (1996), see RN 184920-98-7.	1,2,4,7,8, 12,14-16 ----- 3,5,6,9-11,13,17-86
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:	"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A" document member of the same patent family	
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search 04 MARCH 2003	Date of mailing of the international search report 26 MAR 2003	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3250	Authorized officer CELIA CHANG Telephone No. (703) 305-1435	

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US02/39651

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ----- Y	Database CAS on STN, (Columbus, OH, USA) DN 123:169516, WETTERAU et al. 'Preparation fo acylaminopiperidines and piperazines as inhibitors of microsomal triglyceride transfer protein. (1995) EP 643057, see RN163266-92-0, 163266-94-2.	1,2,4,7,8, 12, 14-16 ----- 3, 5-6, 9-11, 13, 17-41
X ---- Y	Database CAS on STN (Columbus, OH, USA) DN 116:20945, CAIN et al. 'preparation of aryl(alkyl)4-piperidylalkylethers as psychotropics and agrochemical fungicides' (1991) EP 449187, see RN 138030-83-8, 138031-13-7.	1, 2, 4, 7-8, 12, 14-16 ----- 3, 5, 6, 9-11, 13, 17-86
X ---- Y	Database CAS on STN (Columbus, OH, USA) DN 137:279361, GARVEY et al. 'Preparation of nitrosated and hitrosylated alpha-adrenergic receptor antagonists for the treatment of sexual dysfunction' (2002) US 2002143007, see RN 57368-81-7.	1,4,7,8,12, ----- 2-3, 5-6, 9-11, 13- 86
X ---- Y	Database CAS on STN (Columbus, OH, USA) DN 129:302552, KITAZAWA et al. 'Preparation of 1,4-disubstituted cyclic amine derivatives as serotonin antagonists' (1998) WO 9843956, see RN 202859-14-1, 214611-21-9, 131587-28-5.	1,4,7,8,12 ----- 2,3,5,6, 9-11,13-86
X ----- Y	Database CAS on STN (Columbus, OH, USA) DN 111:23496, SEVRIN et al. 'Piperidinylmethyl tetrahydropyridoindoles as drugs with affinity for serotin in receptors and their preparation' (1989) see RN 120848-72-8.	1,4,7,8,12, ----- 2,3,5,6, 9-11,13-86
X ---- Y	Database CAS on STN (Columbus, OH, USA) DN 132:151821, ITO et al. 'Preparation of 2-substituted-1-piperidylbenzimidazoles as ORL1 receptor agonists' (2000) WO0008013, see RN 258289-01-9, 258289-02-0, 231962-49-5, 258287-10-4, 258287-13-7.	1,4,7,8,12 ----- 2,3,5,6, 9-11, 13-86
X ---- Y	US 5,403,847 (gGLUCHOWSKI et al.) 04 April 1995, see entire document especially, figure 1, compound indoromin, 7, 9 and columns 18-19 table 1 and 2.	1,8,9 ----- 2-7, 10-86
X ---- Y	WO 2001.51469 (MERCK PATENT GMBH) 19 July 2001, see entire document, especially, p.31-38 compounds and disorders of the abstract.	1, 8 ----- 2-7, 9-86
X,P	Database CHEMCATS on STN (Columbus, OH, USA) AN 2002:2612714, Interchim Intermediates (2002), see RN 433962-80-2.	1
Y,P	US 2002 0177607 (HOEMANN) 28 November 2002, see entire document, especially claims 1-91.	1-86

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/39651

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

IPC(7): A61K 31/445, 31/454, 31/4545; C07D 211/54, 211/68, 401/02, 401/12